

PATENT COOPERATION TREATY

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Assistant Commissioner for Patents
United States Patent and Trademark
Office
Box PCT
Washington, D.C. 20231
ETATS-UNIS D'AMERIQUE

in its capacity as elected Office

Date of mailing (day/month/year) 15 June 2000 (15.06.00)	
International application No. PCT/US99/18507	Applicant's or agent's file reference
International filing date (day/month/year) 11 August 1999 (11.08.99)	Priority date (day/month/year) 11 August 1998 (11.08.98)
Applicant PAPADOPOULOS, Vassilios et al	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:

23 November 1999 (23.11.99)

☐ in a notice effecting later election filed with the International Bureau on:2. The election ☒ was☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

<p>The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland</p> <p>Facsimile No.: (41-22) 740.14.35</p>	<p>Authorized officer Yolaine CUSSAC</p> <p>Telephone No.: (41-22) 338.83.38</p>
--	--

PATENT COOPERATION TREATY

PCT

INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference	FOR FURTHER ACTION <small>see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.</small>	
International application No. PCT/US 99/ 18507	International filing date (day/month/year) 11/08/1999	(Earliest) Priority Date (day/month/year) 11/08/1998
Applicant GEORGETOWN UNIVERSITY MEDICAL CENTER et al.		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 7 sheets.

☒ It is also accompanied by a copy of each prior art document cited in this report.

1. Basis of the report

- a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.

☐ the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).

- b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing :

☒ contained in the international application in written form.

☐ filed together with the international application in computer readable form.

☐ furnished subsequently to this Authority in written form.

☒ furnished subsequently to this Authority in computer readable form.

☒ the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.

☒ the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☒ **Certain claims were found unsearchable** (See Box I).

3. ☒ **Unity of invention is lacking** (see Box II).

4. With regard to the **title**,

☒ the text is approved as submitted by the applicant.

☐ the text has been established by this Authority to read as follows:

5. With regard to the **abstract**,

☒ the text is approved as submitted by the applicant.

☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the **drawings** to be published with the abstract is Figure No.

☐ as suggested by the applicant.

☐ because the applicant failed to suggest a figure.

☐ because this figure better characterizes the invention.

☒ None of the figures.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 99/18507

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 35-40 -in as far as they concern in vivo methods- are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: 29-32
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

The claimed agents have not been described in the application and have as such not been the subject of search.
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☒ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

1-4, 9-18, 23-25, 27-40 partially and 5, 19, 26 completely
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 29-32

The claimed agents have not been described in the application and have as such not been the subject of search.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-3, 9-18, 24, 25, 27-32, 35-40 all partially

An isolated DNA fragment encoding PBR-associated protein 3 (PAP3) and portions of said DNA, including the DNA of SEQ ID No:1 and fragments of said sequence of at least 30 nucleotides. Recombinant DNA constructs, host cells and production methods relating to PAP3. Recombinant PAP3 and detection methods comprising the two hybrid assay. Antibodies to the PAP3 polypeptide of SEQ ID No:6. Methods for detecting agents which reduce, eliminate or promote PAP3 activity, agents and therapeutic compounds that inhibit or promote PAP activity and therapeutic methods for the treatment of disease states.

2. Claims: 5, 19, 26 completely and 1-4, 9-18, 23-25, 27-40 all partially

As invention 1 but relating to PAP7 (SEQ ID Nos: 2 and 7)

3. Claims: 6, 20 completely and 1-3, 9-18, 23-25, 27-40 all partially

As invention 1 but relating to PAP8 (SEQ ID Nos: 3 and 8)

4. Claims: 7, 21 completely and 1-3, 9-18, 23-25, 27-40 all partially

As invention 1 but relating to PAP15 (SEQ ID Nos: 4 and 9)

5. Claims: 8, 22 completely and 1-4, 9-18, 23, 24, 27-40 all partially

As invention 1 but relating to PAP20 (SEQ ID Nos: 5 and 10)

INTERNATIONAL SEARCH REPORT

International Application No

CT/US 99/18507

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C07K14/47 C12N1/21 G01N33/53 C07K16/18
 C12Q1/68 A61P35/00 A61P3/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K G01N C12Q A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL 'Online! AC AF022770, 1 October 1997 (1997-10-01) LI H & V PAPADOPOULOS: "M. musculus peripheral benzodiazepine receptor associated protein" XP002133903 cited in the application 94.9% identity in 178 aa overlap with SEQ ID No:7 the whole document</p> <p style="text-align: center;">--- -/--</p>	<p>1-5, 9-16, 19</p>



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

15 June 2000

Date of mailing of the international search report

28. 06. 00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Nichogiannopoulou, A

INTERNATIONAL SEARCH REPORT

International Application No

CT/US 99/18507

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL 'Online! AC X64455, 21 February 1992 (1992-02-21) DON J: "M. musculus mRNA megl" XP002133904 cited in the application 99% identity in 490 nt overlap with SEQ ID No:1 (17-505:28-517) the whole document</p> <p>---</p>	3,9-16
X	<p>DATABASE EMBL 'Online! AC AA174581, 31 December 1996 (1996-12-31) MARRA M ET AL: "mo98e12.r1 Stratagene mouse testis (#937308) Mus musculus cDNA" XP002133905 EST with 97.9% identity in 421 nt overlap (17-435:79-498) with SEQ ID No:1 the whole document</p> <p>---</p>	3,9-16
E	<p>WO 00 01821 A (INCYTE PHARMA INC) 13 January 2000 (2000-01-13) Incyte clone with 88.7% identity to SEQ ID NO:2 table 2</p> <p>---</p>	3-5, 9-16,19
P,A	<p>GALIÈGUE S ET AL: "Cloning and characterisation of PRAX-1. A new protein that specifically interacts with the peripheral benzodiazepine receptor" J. BIOL. CHEM, vol. 274, no. 5, 29 January 1999 (1999-01-29), page 2938-2952 XP002122735 cited in the application the whole document</p> <p>---</p>	
A	<p>KRUEGER K E ET AL: "Peripheral-type benzodiazepine receptors mediate translocation of cholesterol from outer to inner mitochondrial membranes in adrenocortical cells" JOURNAL OF BIOLOGICAL CHEMISTRY., vol. 265, no. 25, 5 September 1990 (1990-09-05), pages 15015-15022, XP002133901 ISSN: 0021-9258 cited in the application the whole document</p> <p>---</p> <p style="text-align: center;">-/--</p>	38,39

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/18507

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>MIETTINEN H ET AL: "Expression of peripheral-type benzodiazepine receptor and diazepam binding inhibitor in human astrocytomas: relationship to cell proliferation" CANCER RESEARCH, vol. 55, 15 June 1995 (1995-06-15), pages 2691-2695, XP002133902 cited in the application the whole document -----</p>	36,37

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/18507

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0001821 A	13-01-2000	AU 4857999 A	24-01-2000

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07K 14/00	A2	(11) International Publication Number: WO 00/09549 (43) International Publication Date: 24 February 2000 (24.02.00)
(21) International Application Number: PCT/US99/18507 (22) International Filing Date: 11 August 1999 (11.08.99) (30) Priority Data: 60/096,048 11 August 1998 (11.08.98) US (71) Applicant (for all designated States except US): GEORGETOWN UNIVERSITY MEDICAL CENTER [US/US]; Office of the Dean of Research and Graduate Education, Building D, suite 177, Washington, DC 20007 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): PAPADOPOULOS, Vasilios [US/US]; 15417 Peach Leaf Drive, North Potomac, MD 20878 (US). LI, Hua [CN/US]; Apartment 102, 1319 Fort Myer Drive, Arlington, VA (US). (74) Agent: PRATT, Sana, A.; Pratt & Associates, Inc., 10821 Hillbrooke Lane, Potomac, MD 20854 (US).		(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: PERIPHERAL-TYPE BENZODIAZEPINE RECEPTOR ASSOCIATED PROTEINS, CLONING, EXPRESSION AND METHODS OF USE (57) Abstract The present invention relates to nucleic acids encoding PBR-associated proteins (PAPs) and methods for use in producing PAPs and methods of using PAPs. In this study we have identified proteins (PAPs) that interact with PBR protein by a yeast two-hybrid system. We used PBR as bait to screen a mouse testis cDNA library. Five clones were isolated based on their ability to interact with PBR. These proteins may be involved in the regulation of the function of PBR, serving as an endogenous ligand or allosteric modulator of the receptor.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE OF THE INVENTION

**Peripheral-type Benzodiazepine Receptor Associated
Proteins, Cloning, Expression and Methods of Use**

5

Field of the Invention

The present invention relates to nucleic acid molecules encoding peripheral-type benzodiazepine receptor (PBR) associated proteins (PAPs), including
10 mutants, variants, fragments and derivatives thereof, and to vectors and host cells comprising such nucleic acid molecules; methods of using PAPs; method for screening for inhibitors and activators of PAPs or PBR; and kits comprising the compositions or
15 polypeptides of the invention.

Background of the Invention

The peripheral-type benzodiazepine receptor (PBR) was originally discovered because it binds the
20 benzodiazepine diazepam with relatively high affinity (Papadopoulos, V. 1993, *Endocr. Rev.* 14:222-240). Benzodiazepines are among the most highly prescribed drugs due to their pharmacological actions in relieving anxiety mediated through modulating the
25 activity of γ -aminobutyric acid receptors in the central nervous system (Costa, E. and Guidotti, A. 1979, *Annu. Rev. Pharmacol. Toxicol.* 19:531-545). PBR is another class of binding sites for benzodiazepines distinct from the aforementioned neurotransmitter
30 receptors. Further studies demonstrated that in addition to benzodiazepines, PBR binds other classes of organic compounds with high affinity (Papadopoulos, 1993, *supra*). PBR, although present in all tissues examined, was found to be particularly high in steroid

producing tissues, where it was primarily localized in the outer mitochondrial membrane (OMM) (Anholt, R.R.H. et al. 1986, *J. Biol. Chem.* 261:576-583). An 18 kDa isoquinoline-binding protein was identified as PBR, 5 cloned and expressed (Papadopoulos, V. 1998, *Proc Soc. Exp. Biol. Med.* 217:130-142). It was then demonstrated that PBR is a functional component of the steroidogenic machinery (Papadopoulos, 1998, *supra*; Papadopoulos V. et al. 1990, *J. Biol. Chem.* 265:3772-10 3779) mediating cholesterol delivery from the outer to the inner mitochondrial membrane (Krueger, K. E. and Papadopoulos, V. 1990, *J. Biol. Chem.* 265:15015-15022). Further studies demonstrated that pharmacologically induced reduction of adrenal PBR 15 levels *in vivo* resulted in decreased circulating glucocorticoid levels (Papadopoulos, V. 1998, *supra*). In addition, targeted disruption of the PBR gene in Leydig cells resulted in the arrest of cholesterol transport into mitochondria and steroid formation; 20 transfection of the mutant cells with a PBR cDNA rescued steroidogenesis (Papadopoulos, V. et al. 1997, *J. Biol. Chem.* 272:32129-32135).

PBR is extremely abundant in steroidogenic cells and found primarily on outer mitochondrial membranes 25 (Anholt, R. et al. 1986, *J. Biol. Chem.* 261:576-583). PBR is thought to be associated with a multimeric complex composed of the 18-kDa isoquinoline-binding protein and the 34-kDa pore-forming voltage-dependent anion channel protein, preferentially located on the 30 outer/inner mitochondrial membrane contact sites (McEnery, M. W. et al. *Proc. Natl. Acad. Sci. U.S.A.* 89:3170-3174; Garnier, M. et al. 1994, *Mol. Pharmacol.* 45:201-211; Papadopoulos, V. et al. 1994, *Mol. Cell. Endocr.* 104:R5-R9). Drug ligands of PBR, upon binding 35 to the receptor, simulate steroid synthesis in

steroidogenic cells *in vitro* (Papadopoulos, V. et al. 1990, *J. Biol. Chem.* 265:3772-3779; Ritta, M. N. et al. 1989, *Neuroendocrinology* 49: 262-266; Barnea, E. R. et al. 1989, *Mol. Cell. Endocr.* 64:155-159; Amsterdam, A. and Suh, B. S. 1991, *Endocrinology* 128:503-510; Yanagibashi, K. et al. 1989, *J. Biochem. (Tokyo)* 106: 1026-1029). Likewise, *in vivo* studies showed that high affinity PBR ligands increase steroid plasma levels in hypophysectomized rats (Amri, H. et al. 1996, *Endocrinology* 137:5707-5718). Further *in vitro* studies on isolated mitochondria provided evidence that PBR ligands, drug ligands, or the endogenous PBR ligand, the polypeptide diazepam-binding inhibitor (BDI) (Papadopoulos, V. et al. 1997, *Steroids* 62:21-28), stimulate pregnenolone formation by increasing the rate of cholesterol transfer from the outer to the inner mitochondrial membrane (Krueger, K. E. and Papadopoulos, V. 1990, *J. Biol. Chem.* 265:15015-15022; Yanagibashi, K. et al. 1988, *Endocrinology* 123: 2075-2082; Besman, M. J. et al. 1989, *Proc. Natl. Acad. Sci. U.S.A.* 86: 4897-4901; Papadopoulos, V. et al. 1991, *Endocrinology* 129: 1481-1488).

Based on the amino acid sequence of the 18-kDa PBR, a three dimensional model was developed (Papadopoulos, V. 1996, In: The Leydig Cell. Payne, A. H. et al. (eds) Cache River Press, IL, pp 596-628). This model was shown to accomodate a cholesterol molecule and function as a channel, supporting the role of PBR in cholesterol transport. Recently we demonstrated the role of PBR in steroidogenesis by generating PBR negative cells by homologous recombination (Papadopoulos, V. et al. 1997, *J. Biol. Chem.* 272:32129-32135) that failed to produce steroids. However, addition of the hydrosoluble

analogue of cholesterol, 22R-hydroxycholesterol, recovered steroid production by these cells, indicating that the cholesterol transport mechanism was impaired. Further cholesterol transport
5 experiments in bacteria expressing the 18-kDa PBR protein provided definitive evidence for a function as a cholesterol channel/transporter (Li and Papadopoulos, 1998, *Endocrinology* 139, 4991-4997).

Studies in a number of tumors such as rat brain
10 containing glioma tumors (Richfield, E. K. et al. 1988, *Neurology* 38:1255-1262), colonic adenocarcinoma and ovarian carcinoma (Katz, Y. et al. 1988, *Eur. J. Pharmacol.* 148: 483-484 and Katz, Y. et al. 1990, *Clinical Sci.* 78:155-158) have shown an abundance of
15 peripheral-type benzodiazepine receptors (PBR) compared to normal tissue. All documents cited herein *infra* and *supra* are hereby incorporated in their entirety by reference thereto. Moreover, a 12-fold increase in PBR density relative to normal parenchyma,
20 was found in human brain glioma or astrocytoma (Cornu, P. et al. 1992, *Acta Neurochir.* 119:146-152). The authors suggested that PBR densities may reflect the proliferative activity of the receptor in these tissues. Recently, the involvement of PBR in cell
25 proliferation was further shown (Neary, J. T. et al. 1995, *Brain Research* 675:27-30; Miettinen, H. et al. 1995, *Cancer Research* 55:2691-2695), and its expression in human astrocytic tumors was found to be associated with tumor malignancy and proliferative
30 index (Miettinen, H. et al. *supra*; Alho, H. 1994, *Cell Growth Different.* 5:1005-1014). Characterization of PBR in human breast cancer biopsies, led to the discovery that the invasive and metastatic ability of human breast tumor cells is proportional to the level
35 of PBR expressed, and correlates with the subcellular

localization of PBR in these cells in that PBR is found primarily in the nucleus in aggressive tumor cells whereas PBR is found primarily in the cytoplasm of invasive but non-aggressive cells. These changes
5 in PBR expression can be used as a tool for detection, diagnosis, prevention and treatment in breast cancer patients, in particular, and in aggressive solid tumors in general.

Since both PBR and its endogenous ligand, the
10 polypeptide diazepam binding inhibitor, are constitutively expressed in steroidogenic cells, the regulation of PBR function by hormones may be due to its association with other proteins. This interaction may result in the initiation of steroid biosynthesis.
15 Therefore, there is a need to identify proteins which associate with PBR and may modulate PBR function.

Summary of the Invention

The present invention meets the need described above. We have identified PBR associated proteins (PAPs) that
20 interact with PBR using the two-hybrid system. We used PBR as a bait to screen a mouse testis cDNA library. Five clones were isolated by their ability to interact with PBR: PAP3, PAP7, PAP8, PAP15, and PAP20. Among the nucleotide sequences identified,
25 PAP3 was identical to the previously isolated meg1 protein (Don, J. and Wolgemuth, D. J., 1992, *Cell Growth Differ.* 3, 495; Ever, L. et al., 1999, *Cell Growth Differ.* 10, 19-26). PAP7, PAP8, PAP15, and PAP20 are novel sequences according to a search in the
30 Genebank database which did not find a match for these sequences. PAP7 and PAP17 are different clones of the same novel protein product. All PAPs have fatty acylation (myristoylation) sites and PKC

phosphorylation sites. In addition, PAP20 has a PKA phosphorylation site. The distribution and function of the PAPs, as well as their functional relationship to PBR is under investigation.

- 5 So far, the distribution of PAP7 in major mouse tissues such as brain, testis, ovary, adrenal, kidney and muscle showed a profile similar to the broader expression pattern of PBR, with an expression level paralleling the steroidogenic ability of the tissue.
- 10 These data imply a role for these PAPs in the regulation of PBR function, serving as endogenous ligands or allosteric modulators of the receptor.

 Therefore, it is an object of the present invention to provide novel DNA fragments encoding PBR associated proteins, PAP3 (SEQ ID NO:1), PAP7 (SEQ ID NO:2) and Genbank accession no. AF022770, PAP8 (SEQ ID NO:3), PAP15 (SEQ ID NO:4), and PAP20 (SEQ ID NO:5) GenBank accession no. AF020338. The DNA fragment is useful as a diagnostic agent for detection of nucleic acid sequences encoding PBR-associated proteins, as an agent for preparation of the protein encoded by the DNA, for the preparation of sequences encoding PAPs, whether cDNA or genomic, and as therapeutic agents.

20 It is another object of the invention to provide an amino acid sequence for PAPs encoded by the DNA sequences described above.

25 It is another object of the invention to provide a recombinant vector comprising a vector and the above described DNA fragments.

30 It is a further object of the present invention to provide a host cell transformed with the above-described recombinant DNA construct.

It is another object of the present invention to provide a method for producing PAPs which comprises culturing a host cell under conditions such that the above-described DNA fragment is expressed and a PAP is
5 thereby produced, and isolating the PAP for use as a reagent, for example for screening drugs and inhibitors of PBR or the PAP itself, for diagnosis, and for therapy.

It is a further object of the present invention
10 to provide an antibody to the above-described recombinant PAPs.

It is yet another object of the present invention to provide a method for detecting any of PAP3, PAP7, PAP8, PAP15, or PAP20 in a sample comprising:

- 15 (i) contacting a sample with antibodies which recognize any one of the PAPs mentioned above; and
(ii) detecting the presence or absence of a complex formed between the PAP and antibodies specific therefor.

20 It is a further object of the present invention to provide a diagnostic kit comprising an antibody against PAP and ancillary reagents suitable for use in detecting the presence of PAP in cells, tissue or serum from yeast, mammals, animals, birds, fish, and
25 plants.

It is yet another object of the present invention to provide a method for the detection of PAP from a sample using the polymerase chain reaction.

It is a further object of the present invention
30 to provide a diagnostic kit comprising primers or oligonucleotides specific for PAP RNA or cDNA suitable for hybridization to PAP RNA or cDNA and/or amplification of PAP sequences and ancillary reagents suitable for use in detecting PAP RNA/cDNA in
35 mammalian tissue.

It is yet another object of the present invention to provide a method for the detection of a PAP in a sample which comprises assaying for the presence or absence of PAP RNA or cDNA in a sample by
5 hybridization assays.

It is an object of the present invention to provide a method for the measurement of PBR in a sample. The method comprises measuring the presence of a PAP complexed with PBR.

10 It is yet another object of the present invention to provide a method for modulating the function or altering the targeting of PBR by increasing or decreasing interaction of PBR with a PAP. PBR functions which can be modulated include cholesterol
15 transport into the cell, steroid production, cell proliferation, and embryogenesis.

It is still another object of the present invention to provide a method for increasing or decreasing PBR function or expression in a cell by
20 providing into the cell a PAP such that PBR function is increased or decreased.

It is yet another object of the present invention to provide a method for increasing or reducing steroidogenesis by altering the level of PAP in said
25 cell.

It is still another object of the invention to provide a therapeutic method for the treatment or amelioration of diseases resulting from increased cell proliferation resulting from aberrant PBR function or
30 expression or localization, said method comprising providing to an individual in need of such treatment an effective amount of a PAP, in a pharmaceutically acceptable diluent, of a PAP such that the aberrant PBR expression, function or localization, is
35 corrected.

It is another object of the present invention to provide a therapeutic method for the treatment or amelioration of diseases resulting from decreased cell proliferation, said method comprising providing to an individual in need of such treatment an effective amount of PAP or an antibody against PAP or an agent which inhibits PAP expression or function in a pharmaceutically acceptable excipient.

It is another object of the present invention to provide a therapeutic method for the treatment or amelioration of diseases resulting from decreased or increased steroidogenesis, said method comprising providing to an individual in need of such treatment an effective amount of a PAP or an antibody against PAP or an agent which inhibits or activates PAP expression or function in a pharmaceutically acceptable diluent.

It is yet a further object of the present invention to provide a cDNA sequence encoding PAPs and vectors incorporating all or a fragment of said sequence, and cells, prokaryotic and eukaryotic, transformed or transfected with said vectors, for use in screening agents and drugs which inhibit expression or function of PAPs or PBR in such cells.

BRIEF DESCRIPTION OF THE DRAWINGS

These and other features, aspects, and advantages of the present invention will become better understood with reference to the following description and appended claims, and accompanying drawings where:

Figure 1. PAP7 protein expressed in MA10 mouse Leydig tumor cells. (A) Western blot. (B), Immunocytochemistry.

Figure 2. PAP7 mRNA Tissue distribution analysis by Dot blot analysis. (A), a Master blot

containing 100-500 ng of poly(A)+ RNA from mouse tissues were hybridized at high stringency with a ³²P-labeled PAP7 probe as described under "Materials and Methods." The autoradiogram was exposed overnight.

5 (B), densitometric analysis of PAP7 expression.

Figure 3. PAP7 tissue distribution analysis by Northern blot analysis. (A), Northern blot analysis was performed using 20 µg of total RNA/lane from different mouse tissues as indicated. The blot was
10 hybridized at high stringency with a ³²P-labeled PAP7 probe as described under "Materials and Methods". The autoradiogram was exposed overnight. (B), Densitometric analysis of PAP7 expression.

Figure 4. PBR tissue distribution analysis by Northern blot analysis. (A), Northern blot analysis was performed using 20 µg of total RNA/lane from different mouse tissues as indicated. The blot was
15 hybridized at high stringency with a ³²P-labeled PBR probe as described under "Materials and Methods". The autoradiogram was exposed overnight. (B), For
20 Northern blot analysis, blots were quantitated by densitometry.

Figure 5. Immunohistological staining of mouse tissues with anti-PAP7 antibody.

25 **Figure 6.** The effect of PAP7 on Steroid Biosynthesis. Progesterone formation in MA-10 stimulated by hCG (A), for different time course (B). The results shown represent the means + S.D. from 2 to 6 independent experiments.

30 **Figure 7.** PAP3 mRNA Tissue distribution analysis by Dot blot analysis. (A), a Master blot containing 100-500 ng of poly(A)+ RNA from mouse tissues were hybridized at high stringency with a ³²P-labeled PAP3 probe as described under "Experimental

Procedures." The autoradiogram was exposed overnight.
(B), densitometric analysis of PAP3 expression.

Figure 8. PAP20 mRNA Tissue distribution analysis by Dot blot analysis. (A), a Master blot
5 containing 100-500 ng of poly(A)+ RNA from mouse tissues were hybridized at high stringency with a ³²P-labeled PAP20 probe as described under "Experimental Procedures." The autoradiogram was exposed overnight.
(B), densitometric analysis of PAP20 expression.

10 **DETAILED DESCRIPTION**

The five PAPs described in this application were discovered using a two-hybrid assay. The two-hybrid assay is a yeast-based genetic assay used to detect protein-protein interactions *in vivo*. A positive
15 result obtained with the two-hybrid assay allows rapid identification of genes encoding proteins that interact with a target protein. In addition the two-hybrid assay is a sensitive method for detecting weak and transient interactions, which are probably the
20 norm in large native complexes. Most notably, because the two-hybrid assay is performed *in vivo*, the proteins involved are more likely to be in their native conformations.

The two-hybrid assay is based on the fact that
25 many eukaryotic transcriptional activators consist of two physically separable modular domains: one acts as the DNA-binding domain, while the other functions as the transcriptional activation domain. The DNA-binding domain localizes the transcription factor to
30 specific DNA sequences present in the upstream region of genes that are regulated by this factor, while the activation domain contacts other components of the transcription machinery required to initiate transcription. Both domains are required for normal

activation functioning, and normally the two domains are part of the same protein.

In our PAPs screening experiment, the MATCHMAKER Two-Hybrid System from CLONTECH was used. In the
5 MATCHMAKER System, sequences encoding the two functional domains of the GAL4 transcriptional activator have been cloned into two different shuttle/expression vectors (pGBT9 and pGAD10). The pGBT9 hybrid cloning vector is used to generate a
10 fusion of the GAL4 DNA-binding domain with PBR protein. The pGAD10 hybrid cloning vector is used to generate a fusion of the GAL4 activation domain with a collection of random proteins in a fusion mouse testis library (CLONTECH). Both hybrid proteins are targeted
15 to the yeast nucleus by nuclear localization sequences that either are an intrinsic part of the GAL4 DNA-binding domain or have been added to the activation domain from a heterologous source. If PBR protein and an unknown protein or proteins interact with each
20 other, the DNA-binding domain of GAL4 will be tethered to its transcriptional activation domain, and the proper function of the transcription of an appropriate reporter gene (lacZ or HIS3) containing upstream GAL4 binding sites is used to indicate interaction between
25 the two proteins. This allows a positive selection for clones that are transformed by two interacting hybrid constructs and makes library screening more convenient and practical. After a positive clone has been identified, the gene corresponding to the interacting
30 protein was sequenced using the sequencing primers provided in the kit.

In one embodiment, the present invention relates to a DNA or cDNA sequence encoding PBR associated proteins (PAPs). Five clones were isolated, PAP3,
35 comprising 568 bp and identified in SEQ ID NO:1 (Don,

J. and Wolgemuth, D. J., 1992, *Cell Growth Differ.* 3, 495; Ever, L. et al., 1999, *Cell Growth Differ.* 10, 19-26) which encodes a peptide of 83 amino acids identified in SEQ ID NO:6; PAP7, comprising 577 bp
5 extending from 696 to 1164 of the sequence identified in SEQ ID NO:2, which encodes a polypeptide of 363 amino acids, identified in SEQ ID NO:7; PAP8, comprising 568 bp identified in SEQ ID NO:3, which encode a polypeptide of 190 amino acids, identified in
10 SEQ ID NO:8; PAP15 comprising 490 bp identified in SEQ ID NO:4, which encode a polypeptide of 164 amino acids, identified in SEQ ID NO:9; and PAP20 comprising 588 bp identified in SEQ ID NO:5, which encode a polypeptide of 196 amino acids, identified in SEQ ID
15 NO:10.

PAP3 has been identified as the previously isolated meg 1 protein.

PAP7 and PAP17 are different clones of the same novel protein product. Additional PAP7 sequence has
20 been obtained using the 5', 3'RACE system (CLONTECH) and the near full-length gene is identified in SEQ ID NO:2 including the stop codon and some untranslated sequence at the 3' end. The polypeptide encoded by the DNA sequence would have a calculated molecular
25 weight of about 50 kD. Using an PAP7 antibody produced from the initial isolated DNA fragment of 577 bp, a protein of about 52 kD is immunoprecipitated as shown in the Examples below. Analysis of the protein sequence indicates several concensus sequences and
30 important sites such as: two potential myristoylation sites at positions 262-267 and 271-276 of SEQ ID NO:7 and five PKC phosphorylation sites at 395-396, 113-115, 255-257, 280-282, 331-333, and 339-341 of SEQ ID NO:7, an Acyl-Co-A site at position 24-108 of SEQ ID
35 NO:7, a nuclear localization domain at position 150-

167 of SEQ ID NO:7, a troponin site at position 98-247 of SEQ ID NO:2 and an HSP90 domain at position 126-155 of SEQ ID NO:7. The distribution and expression of PAP7 were examined in major mouse tissues such as
5 brain, testis, ovary, adrenal, and kidney, as well as in tissue culture cell lines such as mouse C6 glioma cells, MA-10 Leydig cells, and Y1 adrenal cortical cells. The PAP7 expression pattern is similar to the broader expression profile of PBR in both tissues and
10 cell lines involved in steroid biosynthesis.

Additionally, both PBR and PAP7 expression level in the cell lines correlated with their steroidogenic biosynthesis ability, which suggests that PAP7 may be involved in steroid biosynthesis through PBR.

15 PAP 8, PAP15, and PAP20, are novel genes. The polypeptide encoded by PAP20 has two potential myristoylation sites, one PKC phosphorylation site and one PKA phosphorylation site. Protein myristoylation enables the protein to attach to the cellular membrane and thus take part in cell signaling (Casey, P.J.
20 1995, *Science* 268, 221-225; Boutin, J. A. 1997, *Cell Signal* 9, 15-35). PAP20 is predominantly expressed in the testis. Interaction of PBR with PAP20 increased the affinity of ligand binding using PK11195 as a
25 ligand. Therefore, it is likely that PAP20 serves to increase or decrease PBR function by modulating PBR's affinity to its endogenous ligand, DBI. The tissue distribution of PAP3 and PAP20 is shown in Figure 7 and Figure 8, respectively.

30 Thus, one aspect of the invention provides an isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence selected from SEQ ID NO:1-9 encoding PAPs polypeptides. It is within the skill of a person with ordinary skill in
35 the art to use the sequences provided herein for the

purpose of cloning cDNA or genomic sequences which
encode other parts or complete portions of the PAP
genes described herein and therefore, these related
sequences are encompassed within the present
5 invention.

In addition, isolated nucleic acid molecules of
the invention include DNA molecules which comprise a
sequence substantially different from those described
above but which, due to the degeneracy of the genetic
10 code, still encode PAPs. Of course, the genetic code
and species-specific codon preferences are well known
in the art. Thus, it would be routine for one skilled
in the art to generate the degenerate variants
described above, for instance, to optimize codon
15 expression for a particular host (e.g., change codons
in the human mRNA to those preferred by a bacterial
host such as *E.coli* or plant host).

Nucleic acid molecules of the present invention
may be in the form of RNA, such as mRNA, or in the
20 form of DNA, including, for instance, cDNA and genomic
DNA obtained by cloning or produced synthetically.
The DNA may be double-stranded or single-stranded.
Single-stranded DNA or RNA may be the coding strand,
also known as the sense strand, or it may be the non-
25 coding strand, also referred to as the antisense
strand.

By "isolated" nucleic acid molecule(s) is
intended a nucleic acid molecule, DNA or RNA, which
has been removed from its native environment. For
30 example, recombinant DNA molecules contained in a
vector are considered isolated for the purposes of the
present invention. Further examples of isolated DNA
molecules include recombinant DNA molecules maintained
in heterologous host cells or purified (partially or
35 substantially) DNA molecules in solution. Isolated

RNA molecules include *in vivo* or *in vitro* RNA transcripts of the DNA molecules of the present invention. Isolated nucleic acid molecules according to the present invention further include such
5 molecules produced synthetically.

The present invention is further directed to nucleic acid molecules encoding portions or fragments of the nucleotide sequences described herein. Fragments include portions of the nucleotide sequences
10 of at least 10 contiguous nucleotides in length selected from any two integers, one of which representing a 5' nucleotide position and a second of which representing a 3' nucleotide position, where the first nucleotide for each nucleotide sequence is
15 position 1. That is, every combination of a 5' and 3' nucleotide position that a fragment at least 10 contiguous nucleotide bases in length or any integer between 10 and the length of an entire nucleotide sequence minus 1.

20 Further, the invention includes polynucleotides comprising fragments specified by size, in nucleotides, rather than by nucleotide positions. The invention includes any fragment size, in contiguous nucleotides, selected from integers between 1- and the
25 entire length of an entire nucleotide sequence minus 1. Preferred sizes include 20-50 nucleotides, 50-300 nucleotides useful as primers and probes. Regions from which typical sequences may be derived include but are not limited to, for example, regions encoding
30 specific epitopes or domains within said sequence, for example, the PBR binding domain extending in SEQ ID NO:1, 2, 3, 4, and 5, potential myristoylation sites at positions 262-267 and 271-276 of SEQ ID NO:7 and five PKC phosphorylation sites at 395-396, 113-115,
35 255-257, 280-282, 331-333, and 339-341 of SEQ ID NO:7,

an Acyl-Co-A site at position 24-108 of SEQ ID NO:7, a nuclear localization domain at position 150-167 of SEQ ID NO:7, a troponin site at position 98-247 of SEQ ID NO:7 and an HSP90 domain at position 126-155 of SEQ ID NO:7, among others.

In another aspect, the invention provides isolated nucleic acid molecules comprising polynucleotides which hybridize under stringent hybridization conditions to a polynucleotide sequence of the present invention described above, or a specified fragment thereof. By "stringent hybridization conditions" is intended overnight incubation at 42°C in a solution comprising: 50% formamide, 5X SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5X Denhardt's solution, 10% dextran sulfate, and 20 g/ml denatured sheared salmon sperm DNA, followed by washing the filters in 0.1X SSC at about 65°C.

The sequences encoding the polypeptides of the present invention or portions thereof may be fused to other sequences which provide additional functions known in the art such as a marker sequence, or a sequence encoding a peptide which facilitates purification of the fused polypeptide, peptides having antigenic determinants known to provide helper T-cell stimulation, peptides encoding sites for post-translational modifications, or amino acid sequences which target the fusion protein to a desired location, e.g. a heterologous leader sequence.

The present invention further relates to variants of the nucleic acid molecules of the present invention, which encode portions, analogs or derivatives of the PAPs of the present invention. A variant may occur naturally, such as a natural allelic variant. By an "allelic variant" is intended one of

several alternate forms of a gene occupying a given locus of a chromosome of an organism. Non-naturally occurring variants may be produced by known mutagenesis techniques. Such variants include those produced by
5 nucleotide substitution, deletion, or addition of one or more nucleotides in the coding or noncoding regions or both. Alterations in the coding regions may produce conservative or nonconservative amino acid substitutions, deletions, or additions. Especially
10 preferred among these are silent substitutions, additions, and deletions which do not alter the properties and activities of PAPs polypeptides disclosed herein or portions thereof. Also preferred in this regard are conservative substitutions.

15 Nucleic acid molecules with at least 90-99% identity to a nucleic acid identified above is another aspect of the present invention. These nucleic acids are included irrespective of whether they encode a polypeptide having PAP activity. By "a polypeptide
20 having PAP activity" is intended polypeptides exhibiting activity similar, but not identical, to an activity of the PAP of the invention, as measured in the assays described below. The biological activity or function of the polypeptides of the present
25 invention are expected to be similar or identical to polypeptides from other organisms that share a high degree of structural identity/similarity.

In another embodiment, the present invention relates to a recombinant DNA molecule that includes a
30 vector and a DNA sequence as described above. The vector can take the form of a plasmid, phage, cosmid, YAC, eukaryotic expression vector such as a DNA vector, *Pichia pastoris*, or a virus vector such as for example, baculovirus vectors, retroviral vectors or
35 adenoviral vectors, and others known in the art. The

cloned gene may optionally be placed under the control of (i.e., operably linked to) certain control sequences such as promoter sequences, or sequences which may be inducible and/or cell type-specific.

5 Suitable promoters will be known to a person with ordinary skill in the art. The expression construct will further contain sites for transcription initiation, termination and, in the transcribed region, a ribosome binding site for translation.

10 Among the vectors preferred for use include pGBT9, PGAD10 (Clontech), PSVzeo (Invitrogen), pBlueScript (Stratagene), pCMV5 (Invitrogen), pCRII (Invitrogen) to name a few.

Introduction of the construct into the host cell
15 can be effected by calcium phosphate transfection, electroporation, infection, and other methods known in the art and described in standard laboratory manuals such as Current Protocols in Molecular Biology, Ausubel, F. M. et al. (Eds), Wiley & Sons, Inc. All
20 documents cited herein supra and infra are hereby incorporated in their entirety by reference thereto.

In a further embodiment, the present invention relates to host cells stably transformed or transfected with the above-described recombinant DNA
25 constructs. The host cell can be prokaryotic (for example, bacterial), lower eukaryotic (for example, yeast or insect) or higher eukaryotic (for example, all mammals, including but not limited to rat and human). Both prokaryotic and eukaryotic host cells
30 may be used for expression of desired coding sequences when appropriate control sequences which are compatible with the designated host are used. Among prokaryotic hosts, *E. coli* is most frequently used. Expression control sequences for prokaryotes include
35 promoters, optionally containing operator portions,

and ribosome binding sites. Transfer vectors compatible with prokaryotic hosts are commonly derived from, for example, pBR322, a plasmid containing operons conferring ampicillin and tetracycline resistance, and the various pUC vectors, which also contain sequences conferring antibiotic resistance markers. These markers may be used to obtain successful transformants by selection. Please see e.g., Maniatis, Fritsch and Sambrook, Molecular Cloning; A Laboratory Manual (1982) or DNA Cloning, Volumes I and II (D. N. Glover ed. 1985) for general cloning methods. The DNA sequence can be present in the vector operably linked to a sequence encoding an IgG molecule, an adjuvant, a carrier, or an agent for aid in purification of PAPs, such as glutathione S-transferase, or a series of histidine residues also known as a histidine tag. The recombinant molecule can be suitable for transfecting plant cells or eukaryotic cells, for example, mammalian cells and yeast cells in culture systems. *Saccharomyces cerevisiae*, *Saccharomyces carlsbergensis*, and *Pichia pastoris* are the most commonly used yeast hosts, and are convenient fungal hosts. Control sequences for yeast vectors are known in the art. Mammalian cell lines available as hosts for expression are known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC), such as HEK293 cells, and NIH 3T3 cells, MA10 Leydig cells, mouse C6 glioma cells, Y1 adrenal cells, and breast cancer cell lines such as MDA-231, MCF-7, to name a few. Suitable promoters are also known in the art and include viral promoters such as that from SV40, Rous sarcoma virus (RSV), adenovirus (ADV), bovine papilloma virus (BPV), and cytomegalovirus

(CMV). Mammalian cells may also require terminator sequences and poly A addition sequences; enhancer sequences which increase expression may also be included, and sequences which cause amplification of the gene may also be desirable. These sequences are known in the art. The transformed or transfected host cells can be used as a source of DNA sequences described above. When the recombinant molecule takes the form of an expression system, the transformed or transfected cells can be used as a source of the protein described below.

In another embodiment, the present invention relates to the PAP polypeptides described above or any allelic variation thereof which is immunologically identifiable with the polypeptides.

A polypeptide or amino acid sequence derived from the amino acid sequences mentioned above, refers to a polypeptide having an amino acid sequence identical to that of a polypeptide encoded in the sequence, or a portion thereof wherein the portion consists of at least 2-5 amino acids, and more preferably at least 8-10 amino acids, and even more preferably at least 11-15 amino acids, or which is immunologically identifiable with a polypeptide encoded in the sequence.

A recombinant or derived polypeptide is not necessarily translated from a designated nucleic acid sequence; it may be generated in any manner, including for example, chemical synthesis, or expression of a recombinant expression system. In addition the polypeptide can be fused to other proteins or polypeptides which increase its antigenicity, such as adjuvants for example.

As noted above, the methods of the present invention are suitable for production of any

polypeptide of any length, via insertion of the above-described nucleic acid molecules or vectors into a host cell and expression of the nucleotide sequence encoding the polypeptide of interest by the host cell.

5 Introduction of the nucleic acid molecules or vectors into a host cell to produce a transformed host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction,

10 infection or other methods. Such methods are described in many standard laboratory manuals, such as Davis et al., Basic Methods In Molecular Biology (1986). Once transformed host cells have been obtained, the cells may be cultivated under any

15 physiologically compatible conditions of pH and temperature, in any suitable nutrient medium containing assimilable sources of carbon, nitrogen and essential minerals that support host cell growth. Recombinant polypeptide-producing cultivation

20 conditions will vary according to the type of vector used to transform the host cells. For example, certain expression vectors comprise regulatory regions which require cell growth at certain temperatures, or addition of certain chemicals or inducing agents to

25 the cell growth medium, to initiate the gene expression resulting in the production of the recombinant polypeptide. Thus, the term "recombinant polypeptide-producing conditions," as used herein, is not meant to be limited to any one set of cultivation

30 conditions. Appropriate culture media and conditions for the above-described host cells and vectors are well-known in the art. Following its production in the host cells, the polypeptide of interest may be isolated by several techniques. To liberate the

35 polypeptide of interest from the host cells, the cells

are lysed or ruptured. This lysis may be accomplished by contacting the cells with a hypotonic solution, by treatment with a cell wall-disrupting enzyme such as lysozyme, by sonication, by treatment with high
5 pressure, or by a combination of the above methods. Other methods of bacterial cell disruption and lysis that are known to one of ordinary skill may also be used.

Following disruption, the polypeptide may be
10 separated from the cellular debris by any technique suitable for separation of particles in complex mixtures. The polypeptide may then be purified by well known isolation techniques. Suitable techniques for purification include, but are not limited to,
15 ammonium sulfate or ethanol precipitation, acid extraction, electrophoresis, immunoadsorption, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction
chromatography, affinity chromatography,
20 immunoaffinity chromatography, size exclusion chromatography, liquid chromatography (LC), high performance LC (HPLC), fast performance LC (FPLC), hydroxylapatite chromatography and lectin chromatography.

25 The recombinant polypeptide or fusion protein can be used, detectably labeled and unlabeled, as a diagnostic tool for the detection of PAPs or for the detection and measurement of PBR. Additionally, these polypeptides can be used in a method for modulating
30 PBR expression. In addition, the recombinant protein can be used as a therapeutic agent to reduce cell death and/or increase cell proliferation via its effect on PBR function. The transformed host cells can be used to analyze the effectiveness of drugs and
35 agents which modulate PBR function, expression or

targeting via their effect on the expression or function of PAPs, such as host proteins or chemically derived agents or other proteins which may interact with the cell to alter the PAP function or expression, thereby modulating PBR function, expression or localization.

In another embodiment, the present invention relates to monoclonal or polyclonal antibodies specific for the above-described recombinant proteins (or polypeptides). For instance, an antibody can be raised against a peptide described above, or against a portion thereof of at least 10 amino acids, preferably, 11-15 amino acids. Persons with ordinary skill in the art using standard methodology can raise monoclonal and polyclonal antibodies to the protein (or polypeptide) of the present invention, or a unique portion thereof. Material and methods for producing antibodies are well known in the art (see for example Goding, in, Monoclonal Antibodies: Principles and Practice, Chapter 4, 1986).

The amount of PAP expression can be detected at several levels. Using standard methodology well known in the art, assays for the detection and quantitation of PAP RNA can be designed, and include northern hybridization assays, *in situ* hybridization assays, and PCR assays, among others. Please see e.g., Maniatis, Fritsch and Sambrook, Molecular Cloning: A Laboratory Manual (1982) or DNA Cloning, Volumes I and II (D. N. Glover ed. 1985), or Current Protocols in Molecular Biology, Ausubel, F. M. et al. (Eds), Wiley & Sons, Inc. for general description of methods for nucleic acid hybridization. Polynucleotide probes for the detection of PAP RNA can be designed from the sequences described in SEQ ID NO: 1-9. For example,

RNA isolated from samples can be coated onto a surface such as a nitrocellulose membrane and prepared for northern hybridization. In the case of *in situ* hybridization of biopsy samples for example, the
5 tissue sample can be prepared for hybridization by standard methods known in the art and hybridized with polynucleotide sequences which specifically recognize PAP RNA. The presence of a hybrid formed between the sample RNA and the polynucleotide can be detected by
10 any method known in the art such as radiochemistry, or immunochemistry, to name a few.

One of skill in the art may find it desirable to prepare probes that are fairly long and/or encompass regions of the amino acid sequence which would have a
15 high degree of redundancy in the corresponding nucleic acid sequences. In other cases, it may be desirable to use two sets of probes simultaneously, each to a different region of the gene. While the exact length of any probe employed is not critical, typical probe
20 sequences are no greater than 500 nucleotides, even more typically they are no greater than 250 nucleotides; they may be no greater than 100 nucleotides, and also may be no greater than 75 nucleotides in length. Longer probe sequences may be
25 necessary to encompass unique polynucleotide regions with differences sufficient to allow related target sequences to be distinguished. For this reason, probes are preferably from about 10 to about 100 nucleotides in length and more preferably from about
30 20 to about 50 nucleotides.

The DNA sequence of PAPs can be used to design primers for use in the detection of PAPs using the polymerase chain reaction (PCR) or reverse transcription PCR (RT-PCR). The primers can
35 specifically bind to the PAP cDNA produced by reverse

transcription of PAP RNA, for the purpose of detecting the presence, absence, or quantifying the amount of PAP by comparison to a standard. The primers can be any length ranging from 7-40 nucleotides, preferably 10-15 nucleotides, most preferably 18-25 nucleotides homologous or complementary to a region of the PAP sequence. Reagents and controls necessary for PCR or RT-PCR reactions are well known in the art. The amplified products can then be analyzed for the presence or absence of PAP sequences, for example by gel fractionation, by radiochemistry, and immunochemical techniques. This method is advantageous since it requires a small number of cells. Once PAP is detected, a determination whether the cell is overexpressing or underexpressing PAP can be made by comparison to the results obtained from a normal cell using the same method. For example, increased PAP7 RNA levels correlate with PBR expression levels, especially in steroidogenic cells, wherein, an increase in steroidogenic capability of the cells correlates with an increase in PBR and PAP7 RNA.

In another embodiment, the present invention relates to a diagnostic kit for the detection of PAP RNA in cells, said kit comprising a package unit having one or more containers of PAP oligonucleotide primers for detection of PAP by PCR or RT-PCR or PAP polynucleotides for the detection of PAP RNA in cells by *in situ* hybridization or northern analysis, and in some kits including containers of various reagents used for the method desired. The kit may also contain one or more of the following items: polymerization enzymes, buffers, instructions, controls, detection labels. Kits may include containers of reagents mixed together in suitable proportions for performing the

methods in accordance with the invention. Reagent containers preferably contain reagents in unit quantities that obviate measuring steps when performing the subject methods.

5 In a further embodiment, the present invention provides a method for identifying and quantifying the level of PAP present in a particular biological sample. Any of a variety of methods which are capable of identifying (or quantifying) the level of PAP in a
10 sample can be used for this purpose.

 Diagnostic assays to detect PAPs may comprise a biopsy or *in situ* assay of cells from an organ or tissue sections, as well as an aspirate of cells from a tumor or normal tissue. In addition, assays may be
15 conducted upon cellular extracts from organs, tissues, cells, urine, or serum or blood or any other body fluid or extract.

 When assaying a biopsy, the assay will comprise, contacting the sample to be assayed with a PAP ligand,
20 natural or synthetic, or an antibody, polyclonal or monoclonal, which recognizes PAP, or antiserum capable of detecting PAP, and detecting the complex formed between PAP present in the sample and the PAP ligand or antibody added.

25 PAP ligands or substrates include for example, PBR, in addition to natural and synthetic classes of ligands and their derivatives which can be derived from natural sources such as animal or plant extracts.

 PAP ligands or anti-PAP antibodies, or fragments
30 of ligand and antibodies capable of detecting PAP may be labeled using any of a variety of labels and methods of labeling for use in diagnosis and prognosis of disease associated with increased cell proliferation, such as cancer, or reduced cell death.
35 Examples of types of labels which can be used in the

present invention include, but are not limited to, enzyme labels, radioisotopic labels, non-radioactive isotopic labels, and chemiluminescent labels.

Examples of suitable enzyme labels include malate
5 dehydrogenase, staphylococcal nuclease, delta-5-steroid isomerase, yeast-alcohol dehydrogenase, alpha-glycerol phosphate dehydrogenase, triose phosphate isomerase, peroxidase, alkaline phosphatase, asparaginase, glucose oxidase, beta-galactosidase,
10 ribonuclease, urease, catalase, glucose-6-phosphate dehydrogenase, glucoamylase, acetylcholine esterase, etc.

Examples of suitable radioisotopic labels include
15 ^3H , ^{111}In , ^{125}I , ^{32}P , ^{35}S , ^{14}C , ^{57}To , ^{58}Co , ^{59}Fe , ^{75}Se , ^{152}Eu , ^{90}Y , ^{67}Cu , ^{21}Ci , ^{211}At , ^{212}Pb , ^{47}Sc , ^{109}Pd , ^{11}C , ^{19}F , ^{123}I , etc.

Examples of suitable non-radioactive isotopic labels include ^{157}Gd , ^{55}Mn , ^{162}Dy , ^{52}Tr , ^{46}Fe , etc.

Examples of suitable fluorescent labels include a
20 ^{152}Eu label, a fluorescein label, an isothiocyanate label, a rhodamine label, a phycoerythrin label, a phycodyanin label, an allophycocyanin label, a fluorescamine label, etc.

Examples of chemiluminescent labels include a
25 luminal label, an isoluminal label, an aromatic acridinium ester label, an imidazole label, an acridinium salt label, an oxalate ester label, a luciferin label, a luciferase label, etc.

Those of ordinary skill in the art will know of
30 other suitable labels which may be employed in accordance with the present invention. The binding of these labels to ligands and to antibodies or fragments thereof can be accomplished using standard techniques commonly known to those of ordinary skill in the art. Typical techniques are described by Kennedy, J. H., et
35 al., 1976 (*Clin. Chim. Acta* 70:1-31), and Schurs, A. H.

W. M., et al. 1977 (*Clin. Chim Acta* 81:1-40).

Coupling techniques mentioned in the latter are the glutaraldehyde method, the periodate method, the dimaleimide method, and others, all of which are
5 incorporated by reference herein.

The detection of antibodies (or fragments of antibodies) of the present invention can be improved through the use of carriers. Well-known carriers include glass, polystyrene, polypropylene,
10 polyethylene, dextran, nylon, amylases, natural and modified celluloses, polyacrylamides, agaroses, and magnetite. The nature of the carrier can be either soluble to some extent or insoluble for the purposes of the present invention. The support material may
15 have virtually any possible structural configuration so long as the coupled molecule is capable of binding to PAP. Thus, the support configuration may be spherical, as in a bead, or cylindrical, as in the inside surface of a test tube, or the external surface
20 of a rod. Alternatively, the surface may be flat such as a sheet, test strip, etc. Those skilled in the art will note many other suitable carriers for binding monoclonal antibody, or will be able to ascertain the same by use of routine experimentation.

25 The ligands or antibodies, or fragments of antibodies or ligands of PAPs discussed above may be used to quantitatively or qualitatively detect the presence of PAP. Such detection may be accomplished using any of a variety of immunoassays known to
30 persons of ordinary skill in the art such as radioimmunoassays, immunometric assays, etc. Using standard methodology well known in the art, a diagnostic assay can be constructed by coating on a surface (i.e. a solid support) for example, a
35 microtitration plate or a membrane (e.g.

nitrocellulose membrane), antibodies specific for PAP or a portion of PAP, and contacting it with a sample from a person suspected of having a PAP related disease. The presence of a resulting complex formed
5 between PAP in the sample and antibodies specific therefor can be detected by any of the known detection methods common in the art such as fluorescent antibody spectroscopy or colorimetry. A good description of a radioimmune assay may be found in Laboratory

10 Techniques and Biochemistry in Molecular Biology. by Work, T.S., et al. North Holland Publishing Company, N.Y. (1978), incorporated by reference herein. Sandwich assays are described by Wide at pages 199-206 of Radioimmune Assay Method, edited by Kirkham and
15 Hunter, E. & S. Livingstone, Edinburgh, 1970.

The diagnostic methods of this invention can be predictive of diseases involving PBR including gallstones, atherosclerosis, Niemann-Pick C, Sitosterolemia, Dystrophy, Tumor proliferation
20 (tumorigenesis), Schnyder's corneal crystalline dystrophy. Brain disorders include cholesterol metabolism and Alzheimer's disease, Tellurium toxicity, Smith-Lemli-Opitz syndrome, myelinization, developmental abnormalities and demyelization :
25 Charcot-Marie-Tooth disease; Pelizaeus-Merzbacher disease, Multiple sclerosis, SLA, to name a few. Alternatively, the methods and compositions may be useful as prophylactic treatment, or in screening for compounds effective in prophylactic treatments.

30 The recombinant protein can be used to identify inhibitors or activators of a PAP activity which allows the identification of drugs or agents which modulate PBR activity. Using an assay as described below in the Examples, or for example, introducing a

drug or agent to a cell expressing a PAP and detecting a increase or decrease in the level of PAP RNA or protein, natural and synthetic agents and drugs can be discovered which result in a reduction or elimination, or increase in a PAP activity. Knowledge of the mechanism of action of the inhibitor or activator is not necessary as long as a decrease or increase in the activity of a PAP is detected. Inhibitors may include agents or drugs which either bind or sequester the PAP's substrate(s), such as PBR, or cofactor(s), or inhibit PAP itself, directly, for example by irreversible binding of the agent or drug to the PAP, or indirectly, for example by introducing an agent which binds the competes with PAP binding to its substrate. Activators may include cofactors necessary for proper PAP function or agents which allow a higher turnover rate of binding or release of the PAP to/from PBR or the particular PAP substrate. Agents or drugs related to this invention may result in partial or complete inhibition or various degrees of activation of PAP which may or may not result in modulation of PBR function. Inhibitors or activators of PAP activity may be used in the treatment or amelioration of conditions such as stress, cancer, neurodegenerative disorders, i.e. stroke, Alzheimer's, developmental disorders, infertility, and immune disorders.

Agents which decrease the level of PAP (i.e. in a human or an animal) or reduce or inhibit PAP activity may be used in the therapy of any disease associated with the elevated levels of PAP. Similarly, agents which increase the level of PAP or activate PAP activity may be used in the therapy of any disease associated with reduced levels of PAP. An increase or decrease in the level of PAP is determined when the

change in the level of PAP is about 2-3 fold higher or lower than the level of PAP in the normal cell, up to about 10-100 fold higher or lower than the amount of PAP in a normal cell. Agents which decrease PAP RNA include, but are not limited to, one or more ribozymes capable of digesting PAP RNA, or antisense oligonucleotides capable of hybridizing to PAP RNA such that the translation of PAP is inhibited or reduced resulting in a decrease in the level of PAP. These agents can be administered as DNA, as DNA entrapped in proteoliposomes containing viral envelope receptor proteins (Kanoda, Y. et al., 1989, *Science* 243, 375) or as part of a vector which can be expressed in the target cell such that the DNA or RNA is made. Vectors which are expressed in particular cell types are known in the art, for example, for the mammary gland, please see Furth, (1997) (*J. Mammary Gland Biol. Neopl.* 2, 373) for examples of conditional control of gene expression in the mammary gland. Alternatively, the DNA can be injected along with a carrier. A carrier can be a protein such as a cytokine, for example interleukin 2, or polylysine-glycoprotein carrier. Such carrier proteins and vectors and methods of using same are known in the art. In addition, the DNA could be coated onto tiny gold beads and said beads introduced into the skin with, for example, a gene gun (Ulmer, J. B. et al., 1993, *Science* 259, 1745).

Alternatively, antibodies, or compounds capable of reducing or inhibiting PAP activity, that is reducing or inhibiting either the expression, production or activity of PAP, such as antagonists, can be provided as an isolated and substantially purified protein, or as part of an expression vector capable of being expressed in the target cell such

that the PAP-reducing or inhibiting agent is produced. Similarly, compounds capable of increasing or activating PAP activity, that is increasing or activating either the expression, production, or activity of PAP, such as agonists, can be provided as an isolated and substantially purified protein, or as part of an expression vector capable of being expressed in the target cell such that the PAP-elevating or activating agent is produced. In addition, factors which affect the stability of the protein, and co-factors such as various ions, i.e. Ca^{2+} (Calvo, D. J. and Medina, J. H., 1993, *J. Recept. Res.* 13:975-987), or anions, such as halides or anion channel blockers such as DIDS (4,4'-diisothiocyanostilbene-2,2'-disulfonic acid), an ion transport blocker (Skolnick, P., 1987, *Eur. J. Pharmacol.* 133:205-214), or factors which affect the stability of PBR such as lipids, for example, the phospholipids phosphatidylserine and phosphatidylinositol whereby the presence of the phospholipids is required for receptor activity (Moynagh, P. N. and Williams, D.C., 1992, *Biochem. Pharmacol.* 43:1939-1945) can be administered to modulate the expression and function of the PAPs and PBR. These formulations can be administered by standard routes. In general, the combinations may be administered by the topical, transdermal, intraperitoneal, oral, rectal, or parenteral (e.g. intravenous, subcutaneous, or intramuscular) route. In addition, PAP-inhibiting or PAP-activating compounds may be incorporated into biodegradable polymers being implanted in the vicinity of where drug delivery is desired, for example, at the site of a tumor or implanted so that the PAP-inhibiting or PAP-activating compound is slowly released systemically.

The biodegradable polymers and their use are described, for example, in detail in Brem et al. (1991) *J. Neurosurg.* 74, 441-446. These compounds are intended to be provided to recipient subjects in an amount sufficient to effect the inhibition of PAP. Similarly, agents which are capable of negatively or positively effecting the expression, production, stability or function of PAP, are intended to be provided to recipient subjects in an amount sufficient to result in the desired effect. An amount is said to be sufficient to "effect" the inhibition or induction of PAP if the dosage, route of administration, etc. of the agent are sufficient to influence such a response.

The PAPs identified in this application were discovered due to their ability to associate with PBR, and may play a role in the proper targeting, function, expression, or stability of PBR. Therefore, a method for inhibiting or reducing PBR function, or altering the localization of PBR, would include a method for dissociating PAPs from the receptor. This is possible using agents which block the site on PBR at which these PAPs associate with PBR, or alternatively, blocking the site on the PAPs which is involved in PBR-association. Such agents would include antibodies or antagonists which recognize such sites or which alter the conformation of these sites such that PAP and PBR association is inhibited or eliminated. Agents which decrease the level of PBR (i.e. in a human or an animal) or reduce or inhibit PBR activity may be used in the therapy of any disease associated with the elevated levels of PBR such as metastatic cancer, for example breast cancer, or diseases associated with increased cell proliferation or increased cholesterol transport into the cell. An increase in the level of PBR is determined when the

level of PBR in a tumor cell is about 2-3 times the level of PBR in the normal cell, up to about 10-100 times the amount of PBR in a normal cell.

Antibodies or compounds capable of reducing or
5 inhibiting the association between PBR and PAPs can be provided as an isolated and substantially purified protein, or as part of an expression vector capable of being expressed in the target cell such that the PBR-PAP-association reducing or inhibiting agent is
10 produced. These formulations can be administered by standard routes. In general, the combinations may be administered by the topical, transdermal, intraperitoneal, oral, rectal, or parenteral (e.g. intravenous, subcutaneous, or intramuscular) route.
15 In addition, these compounds may be incorporated into biodegradable polymers being implanted in the vicinity of where drug delivery is desired, for example, at the site of a tumor or implanted so that the compound is slowly released systemically. The biodegradable
20 polymers and their use are described, for example, in detail in Brem et al. (1991) *J. Neurosurg.* 74, 441-446.

These compounds are intended to be provided to recipient subjects in an amount sufficient to effect the inhibition of PBR/PAP association.

25 In line with the function of PBR in cell proliferation, agents which stimulate the function of PBR by increasing the association of PAPs to PBR, may be used in the therapy of any disease associated with a decrease of PBR, or a decrease in cell
30 proliferation, wherein PBR is capable of increasing such proliferation, e.g. developmental retardation. PBR has also been shown to be involved in cholesterol transport, therefore, an agent or drug which results in an increase in function or stability of PBR and its
35 associated PAPs can be used to increase cholesterol

transport into cells. Diseases where cholesterol transport is deficient include lipoidal adrenal hyperplasia, and diseases where there is a requirement for increased production of compounds requiring
5 cholesterol such as myelin and myelination including Alzheimer's disease, spinal chord injury, and brain development neuropathy [Snipes, G. and Suter, U. (1997) Cholesterol and Myelin. In: Subcellular Biochemistry, Robert Bittman (ed.), vol. 28, pp.173-
10 204, Plenum Press, New York], to name a few.

In providing a patient with any agent which modulates the expression, function, targeting, or association of PAP or PBR as discussed above, the dosage of administered agent will vary depending upon
15 such factors as the patient's age, weight, height, sex, general medical condition, previous medical history, etc. In general, it is desirable to provide the recipient with a dosage of agent which is in the range of from about 1 pg/kg to 10 mg/kg (body weight
20 of patient), although a lower or higher dosage may be administered.

A composition is said to be "pharmacologically acceptable" if its administration can be tolerated by a recipient patient. Such an agent is said to be
25 administered in a "therapeutically effective amount" if the amount administered is physiologically significant. An agent is physiologically significant if its presence results in a detectable change in the physiology of a recipient patient.

30 The compounds of the present invention can be formulated according to known methods to prepare pharmaceutically useful compositions, whereby these materials, or their functional derivatives, are combined in admixture with a pharmaceutically

acceptable carrier vehicle. Suitable vehicles and their formulation, inclusive of other human proteins, e.g., human serum albumin, are described, for example, in Remington's Pharmaceutical Sciences [16th ed.,

5 Osol, A. ed., Mack Easton PA. (1980)]. In order to form a pharmaceutically acceptable composition suitable for effective administration, such compositions will contain an effective amount of the above-described compounds together with a suitable
10 amount of carrier vehicle.

Additional pharmaceutical methods may be employed to control the duration of action. Control release preparations may be achieved through the use of polymers to complex or absorb the compounds. The
15 controlled delivery may be exercised by selecting appropriate macromolecules (for example polyesters, polyamino acids, polyvinyl, pyrrolidone, ethylenevinylacetate, methylcellulose, carboxymethylcellulose, or protamine sulfate) and the
20 concentration of macromolecules as well as the method of incorporation in order to control release. Another possible method to control the duration of action by controlled release preparations is to incorporate the compounds of the present invention into particles of a
25 polymeric material such as polyesters, polyamino acids, hydrogels, poly(lactic acid) or ethylene vinylacetate copolymers. Alternatively, instead of incorporating these agents into polymeric particles, it is possible to entrap these materials in
30 microcapsules prepared, for example, interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly(methylmethacrylate)microcapsules, respectively, or in colloidal drug delivery systems, for example,
35 liposomes, albumin microspheres, microemulsions,

nanoparticles, and nanocapsules or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences (1980).

5 The present invention also provides kits for use in the diagnostic or therapeutic methods described above. Kits according to this aspect of the invention may comprise one or more containers, such as vials, tubes, ampules, bottles and the like, which may comprise one or more of the compositions of the
10 invention.

The kits of the invention may comprise one or more of the following components, one or more compounds or compositions of the invention, and one or more excipient, diluent, or adjuvant.

15 It will be readily apparent to one of ordinary skill in the relevant arts that other suitable modifications and adaptations to the methods and applications described herein are obvious and may be made without departing from the scope of the invention
20 or any embodiment thereof. Having now described the present invention in detail, the same will be more clearly understood by reference to the following examples, which are included herewith for purposes of illustration only and are not intended to be limiting
25 of the invention.

The following Materials and Methods were used in the Examples described below.

Material and Methods

Materials

30 [α -³²P]dCTP (specific activity, 3000 Ci/mmol), [1,2,6,7-N-³H]progesterone (specific activity, 94.1 Ci/mmol) and ³H-1-(2-chlorophenyl)-N-methyl-N-(1-methyl-propyl)-3-isoquinolinecarboxamide (PK 11195) (specific activity, 86.9 Ci/mmol) were obtained from

NEN Life Science Products (Boston, MA). PK11195 was obtained from Research Biochemicals, Inc. (Natick, MA). Nitrocellulose (0.45 μ m) was from Hoeffer Scientific (San Francisco, CA). 22R

- 5 Hydroxycholesterol was purchased from Sigma. Restriction enzymes were from Stratagene (La Jolla, CA) and New England Biolabs (Beverly, MA). Cell culture supplies were purchased from from Life Technologies, Inc. (Grand Island, NY). Tissue culture
10 plasticware was from Corning (Corning, NY). Electrophoresis reagents and materials were supplied from BioRad. All other chemicals used were of analytical grade and were obtained from various commercial sources.

15 Strains and media

- The genotype of the *Saccharomyces cerevisiae* reporter strain HF7c is *MATa*, *ura3-52*, *his3-200*, *lys2-801*, *ade2-101*, *trp1-901*, *leu2-3, 112*, *gal4-542*, *gal80-538*, *LYS2::GAL-HIS3*, *URA3::(GAL4 17-mers)₃-CYC1-lacZ*
20 (CLONTECH, Palo Alto, CA). Yeast strains were grown at 30°C in standard liquid YPD medium or minimal SD synthetic medium with appropriate supplement amino acids (CLONTECH, Palo Alto, CA).

Plasmids and construction

- 25 The mouse PBR cDNA coding sequence was subcloned into pGBT9 (CLONTECH, Palo Alto, CA) at *EcoR I* and *BamH I* sites (pGBT-PBR). The fusion site was verified by sequencing. Functional fusion PBR protein, expressed in yeast cells, was verified by PBR ligand
30 binding assay. Mouse testis cDNA library was constructed in pGAD10 [LEU2, GAL4 (768-881)] (CLONTECH, Palo Alto, CA). Amplification of premade libraries was performed by growing the transformants on LB-agar-ampicillin and purifying the plasmids DNA

with QIAGEN Plasmid Giga kit (QIAGEN, Valencia, CA).
In the transfection experiments, PAP7 partial sequence
(including 192 Amino acids C-terminal sequence) was
inserted into pSVzeo vector (Invitrogen, Carlsbad ,
5 CA) at *EcoRI* and *BamHI* sites.

Yeast two-hybrid screening

The Clontech MATCHMAKER two-hybrid system was
applied in this study (detailed in manufacturer's
instruction book). Briefly, the yeast reporter host
10 strain HF7c was simultaneously cotransformed with both
pGBT-PBR and the mouse testis cDNA library in pGAD10
plasmid by using lithium acetate high-efficiency
method (Gietz, D. et al., 1992, *Nucleic Acids Res.* 20,
1425). HIS positive clones were further selected by
15 colony lift filter assay for β -galactosidase activity.
Plasmid DNA was rescued in *Escherichia coli* DH5 α from
yeast cells. Plasmids were retransformed into yeast
HF7c cells with plasmid pGBT-PBR to test for histidine
prototrophy and β -galactosidase activity (Clontech
20 manual). The cDNA inserts from the positive clones
were sequenced. The full length PAP7 cDNA was
obtained by using 5' and 3' RACE kit from kit
(CLONTECH, Palo Alto, CA).

Sequence analysis

25 The ABI PRISMTM dyes terminator cycle sequencing
ready reaction kit (PE Biosystems, Foster City , CA)
and an Applied Biosystems sequencer were used for
sequencing (Applied Biosystems, Foster City , CA) at
the Lombardi Cancer Center Sequencing Core Facility
30 (Georgetown University). DNA sequences were analyzed
by using Entrez and BLAST program against GeneBankTM
Database.

Cell culture transient transfection

MA-10 cells were grown in modified Waymouth's MB752/1 medium containing 15% horse serum, as described previously (Papadopoulos, V. et al., 1990, *J. Biol. Chem.* 265, 3772-3779). Mouse C6 glioma and mouse Y1 adrenal cortical cells were cultured in DMEM and DMEM F12 respectively, with 10% fetal bovine serum. MA10 cells were transiently transfected by electroporation (El Hefnawy, T. et al., 1996, *Mol. Cell Endocrinol.* 119, 207-217). Each Genepulser cuvette (0.4 cm-gap, BioRad, Hercules, CA) contained 8×10^6 cells in 350 μ l antibiotic-free complete Waymouth's growth medium (see above), plus 30 μ g plasmids DNA in 50 μ l of 0.1 x TE. Cells in electroporation cuvettes were electro-shocked at 330 V and at a capacitance of 950 μ Fd generated from Genepulser (BioRad, Hercules, CA). The cells were kept immediately on ice for 10 min before plated into 96 well plates.

Radioligand binding assays

^3H -1-(2-chlorophenyl)-N-methyl-N-(1-methyl-propyl)-3-isoquinolinecarboxamide (PK 11195) (NEN, Boston, MA) binding studies were performed as we previously described (Papadopoulos, V. et al., 1990, *supra*; Garnier, M. et al., 1994, *Molecular Pharmacology* 45, 201-211). The dissociation constant (Kd) and the number of binding sites (Bmax) were determined by Scatchard plot analysis of the data using the LIGAND program (Munson, P. J. and Rodbard, D., 1980, *Anal. Biochem.* 107, 220-239).

RNA (Northern) blot analysis

Total tissue and cellular RNA was isolated by the acid guanidinium thiocyanate-phenol-chloroform extraction method using RNA STAT60 reagent (Tel-Test

Inc., Friendswood, TX). RNA was separated by denature electrophoresis and transferred to Nytran membrane (Schleicher & Schuell Inc., Keene, NH). The RNA blots were hybridized with ^{32}P labeled PAP7 cDNA probe
5 generated from random priming (Boehringer Mannheim, Indianapolis, IN). Autoradiography was performed by exposing Kodak X-Omat AR films (Eastman Kodak, Rochester, NY) to the blots at -80°C overnight.

Steroid biosynthesis

10 MA-10 cells were plated into 96-well plate at the density of 2.5×10^4 /well for overnight. The cells were stimulated with 50 ng/ml hCG in 0.2 ml/well serum-free medium for 2 hours. The culture medium was collected and tested for progesterone production by RIA. The
15 assay was carried out by using anti-progesterone antisera (ICN, Costa Mesa, CA), following the conditions recommended by the manufacturer. The progesterone production was normalized by the amount of protein in each well. Radioimmunoassay data was
20 analyzed using the software provided by Wallac (EG&G Wallac, Gaithersburg, MD).

Antibody generation and Western analysis

Rabbit anti-PAP7 antibody was prepared by sequential immunization with a peptide
25 SSDEEEEEENVTCEEKAKKNANKP (SEQ ID NO:11) of PAP7 protein, which was coupled to KLH. PAP7 antibodies were purified by an affinity resin containing the same peptide immobilized onto agarose (Bethyl Laboratories, Montgomery, TX). MA10 cells were solubilized in
30 sample buffer (25 mM Tris-HCl (pH6.8), 1% SDS, 5% β -mercaptoethanol, 1 mM EDTA, 4% glycerol, and 0.01% bromophenol blue), boiled for 5 min, and loaded onto a 15% SDS-PAGE minigel (MiniProtein II System, BioRad, Richmond, CA). Electrophoresis was performed at 25

mA/gel using a standard SDS-PAGE running buffer (25 mM Tris, 192 mM glycine, and 0.1% SDS). The proteins were electrophoretically transferred to nitrocellulose membrane (Schleicher & Schuell Inc., Keene, NH). The membrane was incubated in blocking buffer (TTBS buffer (20 mM Tris/HCl, pH 7.5, 0.5M NaCl, and 0.05% Tween-20) containing 10% Carnation nonfat milk) at room temperature for 1 hour, followed by incubation with a primary antibody against PAP7 (1:2000) for 2 hours. The membrane was washed with TTBS three times for 10 min each time. After one-hour incubation with the secondary antibody, goat anti-rabbit IgG conjugated with HRP (Signal Transduction), the membrane was washed with TTBS three times for 10 min each time. Specific protein bands were detected by chemiluminescence using the Renaissance Kit (DuPont-New England Nuclear, Wilmington, DE) according to manufacturer's directions.

Immunocytochemistry

MA-10 cells were cultured on four-chambered SuperCell Culture Slides (Fisher Scientific, Pittsburgh, PA) and fixed with methanol at 4°C for 15 min. The fixed cells were incubated with PAP7 antibody (1:250 dilution) with or without PAP7 peptide for 1 hour. After washing, the cells were incubated with HRP conjugated goat anti-rabbit secondary antibody (Transduction Lab, Lexington, KY) for 1 hour. PAP7 staining was visualized with peroxidase using AEC (3-amino-9-ethyl carbazole) as a chromogen to yield a red reaction product. After counterstaining with hematoxylin, slides were dehydrated and permanently mounted.

Immunohistochemistry

Mouse tissues were freshly snapped in liquid nitrogen. Specimens were fixed in cold methanol right after sectioning for 5 min. The slides were then placed in a chamber containing 0.3% H₂O₂ solution in methanol for 20 min at room temperature to inhibit the endogenous peroxidase activity and then incubated in blocking solution (10% goat serum) (Zymed, South San Francisco, CA) for 15 min. Subsequently, the slides were incubated with anti-PAP7 antibody (1:250) for 2 h at room temperature, washed with water and PBS, incubated with HRP conjugated goat anti-rabbit secondary antibody for 1 hour at room temperature, and then washed with PBS. After treatment with AEC reagent for 1 hr at 37°C for color staining, the sections were counterstained with hematoxylin, dehydrated and permanently mounted.

Protein quantification and Statistical analysis

Proteins were quantified by dye-binding assay of Bradford (Bradford, M. M., 1976, *Anal. Biochem.* 72, 248-254) with bovine serum albumin as the standard. Statistical analysis was performed by ANOVA followed by the Student-Newman-Keuls test or the Dunnett multiple comparisons test using the Instat (v.2.04) package from GraphPad, Inc. (San Diego, CA).

Example 1

Isolation of PBR Associated Proteins

We have used the MATCHMAKER Two-Hybrid System from CLONTECH in order to clone genes whose products interact with PBR protein. GAL4 (1-147)-PBR fusion (plasmid pGBT9 + PBR) was used as a bait to screen a mouse MATCHMAKER testis cDNA library constructed into the pGAD10 two-hybrid vector. About 3x10⁶ transformants were tested, and five positive clones were obtained for their ability to interact with PBR.

Library plasmids from these transformants were rescued in *E. coli* strain DH5 α . Both the His⁺ phenotype and the expression of β -galactosidase were confirmed by a second-round transformation of strain HF7c carrying pGBT9-PBR (Table 1).

Table 1: Summary of yeast two hybrid screen of mouse testis library by PBR

Clone	His3	β -Galactosidase
activity		
PAP3	+	++
PAP7	+	++
PAP20	+	++
Positive control	+	+++

Plasmids from these positive clones were first analyzed by restriction enzyme digestion and followed by sequence analysis. Two clones were shown to be coded by single gene, which was an unknown gene and was named as PBR associated protein 7 (PAP7). The other three clones encoded different products. After 5' RACE and 3' RACE, PAP7 cDNA clone was completely sequenced on both strands (SEQ ID NO:2) and it encoded a 463-amino acid protein with a calculated molecular weight of about 52 kDa. A homology search in the Genebank database using the BLAST program showed that this is a new sequence, previously unidentified.

Example 2

PAP7 protein expression in MA-10 Leydig tumor cells

The total MA-10 cell protein extracts were analyzed by western blot using PAP7 antibody. This antibody specifically recognizes a 50 kDa-protein band (Fig. 1A). The PAP7 protein expression in MA-10 cell

was also checked by immunocytochemistry. PAP7 antibody specifically stained MA-10 cell, with the signal mostly localized in the cytoplasm (Figure 1B). This signal can be neutralized by PAP7 peptide, which
5 was used to generate and purify this antibody.

Example 3

PAP7 cells and tissue expression by Dot and Northern Blot

By dot blot analysis, PAP7 was observed to be
10 highly expressed in brain, eye, submax gland, testis, and ovary. Interestingly, PAP7 expression was at its highest level at early embryonic stage, and decreased before birth (Fig. 2A and 2B). Consistently, PAP7 mRNA was expressed in adrenal, brain, heart, liver, testis
15 and ovarian tissues by Northern blot analysis. PAP7 had a 1 kb transcript which was only expressed in testis and a 3 Kb major mRNA transcript in the other tissues (Fig. 3A and 3B). PAP7 was also highly expressed in three cell lines, C6 glioma, MA-10 Leydig
20 cells and Y1 adrenal cells, which have been widely used for studying steroid biosynthesis. All three cell lines expressed PAP7 transcript of the same molecular weight size as in normal tissues. The PAP7 expression level in these cell lines was
25 proportionally correlated with their steroidogenic capability (Fig. 3A and 3B). The PBR mRNA expression level was also checked in these same tissues and cell lines. The PBR expression level was parallel with PAP7 mRNA expression pattern, especially in those
30 three cell lines (Fig. 4A and 4B).

Example 4

PAP7 Cellular distribution

PAP7 protein expression in different tissues was checked by immunohistochemistry (Fig 5). PAP7 was

present in both Leydig and germ cells in testis (Figure 5C and 5D), in hippocampus and neuronal cells in brain (Figure 5E and 5F), in fasciculata cells in adrenal gland (Figure 5G and 5H), and in granulosa cells in ovary (Figure 5A and 5B). Liver and kidney expressed low level of PAP7 protein (data not shown). Each specimen was stained with PAP7 peptide neutralized antibody as a negative control. Subsequent in situ hybridization studies showed that PAP7 mRNA followed PAP-7 protein expression.

Example 5

The effect of PAP7 on steroid biosynthesis in MA-10 cell

PAP7 partial sequence including PBR binding domain was subcloned into pSVzeo mammalian expression vector. This pSVzeoPAP7 vector was transiently transfected into MA-10 cells. pSVzeo empty vector was also transfected into cells as control. The capability of steroid biosynthesis of both empty vector pSVzeo transfectants and pSVzeoPAP7 transfectants was checked by monitoring the progesterone production in response to hormonal (hCG) stimulation. PAP7 transfectants had significantly reduced the level of progesterone production in MA-10 cells as compared with pSVzeo vector transfectant at a dose and time dependent manner (Fig 6).

Discussion

In order to better understand the mechanism of how PBR regulates cholesterol transport activity in steroid biosynthesis, we performed yeast two-hybrid assay to identify the PBR associated protein(s). Mouse PBR cDNA was inserted into the pGBT9 vector to generate a GAL4 DNA-binding domain and PBR fusion

protein as bait. The receptor ligand binding activity of the fusion protein with PBR ligand, PK11195, was tested. Our result indicated that the fusion PBR protein expressed in yeast possessed the similar
5 binding affinity as the native PBR protein (data not shown). PBR has been identified in various peripheral tissues (Gavish, M. and Weizman, R., 1997, *Clin. Neuropharmacol.* 20, 473-481) including testis. Testis is one of the important and very well studied tissues
10 for steroidogenesis (Huhtaniemi, I, and Toppari, J., 1995, *Adv. Exp. Med. Biol.* 377, 33-54). The steroid biosynthesis in the mouse testicular Leydig cell and the role of PBR in this process are also well documented (Papadopoulos, V. et al., 1997, *Steroids* 62,
15 21-28; Papadopoulos, V. et al., 1998, *Endocr. Res.* 24, 479-487). Therefore, we applied mouse testis cDNA library in this two-hybrid screen study. The pGAD10 vector was used to generate a fusion protein of the GAL4 activation domain with a collection of random
20 proteins in the fusion Balb/c mouse testis library. Through the yeast two hybrid screen, PAP7 was identified as one of the positive clones, which demonstrated its ability to interact with PBR (Table 1). Thus, we cloned the PAP7 cDNA coding for a mouse
25 protein that interacted with PBR. Based on the database search, PAP7 is a novel gene product. Recently, PRAX-1 was reported as a new protein that specifically interacts with PBR (Galiege, S. et al., 1999, *J. Biol. Chem.* 274, 2938-2952). The only
30 similarity is that both proteins contain glutamic-acid stretches. Part of PAP7 shares quite high homology with a *C. elegans* gene that has an unknown function (Wilson, R. et al., 1994, *Nature* 368, 32-38). In fact, cholesterol is required for *C. elegans* cell
35 culture (Brenner, S., 1974, *Genetics* 77, 71-94).

Considering that PBR is involved in cholesterol transport and PBR gene is highly conservative in all type of organisms, this data suggests that PAP7 expression may be needed to meet basic requirements
5 for cell survival and growth. PAP7 also shares some homologies with RALBP, a hydrophobic ligand-binding protein that functions in intracellular retinoid transport (Ozaki, K. et al., 1994, *J. Biol. Chem.* 269, 3838-3845).

10 By sequence motif analysis using Swiss-Port Prosite profile scan, PAP7 has fatty acylation (myristoylation) sites, Acyl-CoA-binding protein signature and PKC phosphorylation sites. Protein myristoylation enables protein to attach to the
15 cellular membrane and thus take part in cell signaling (Casey, P.J., 1995, *Science* 268, 221-225; Boutin, J. A., 1997, *Cell Signal* 9, 15-35). PBR is a hydrophobic protein and tightly associated with the outer mitochondrial membrane. This property could enable
20 PAP7 passing hormone stimulation signal to and interacting with PBR thus regulating PBR activity in cholesterol transport. Interestingly, Acyl-CoA-binding protein is the other name of PBR endogenous ligand, diazepam binding inhibitor (DBI) (Rose, T. M.
25 et al., 1992, *Proc. Natl. Acad. Sci. U.S.A.* 89, 11287-11291; Costa, E. and Guidotti, A., 1991, *Life Sci.* 49, 325-344; Suk, K. et al., 1999, *Biochim. Biophys. Acta* 1454, 126-131). This information suggests that PAP7 may coordinate with other PBR endogenous ligands to
30 fulfill its function. The fact that PAP7 has the potential protein kinase phosphorylation sites raises another possibility that PBR could be regulated by hormone stimulation through the interaction with PAP7 protein.

The distribution and expression of PAP7 were examined in major mouse tissues such as brain, testis, ovary, adrenal, and kidney, as well as some cell lines. The PAP7 expressing pattern is similar to the broader expression profile of PBR. According to the previous studies (Papadopoulos, V. et al., 1998, supra), glucocorticoids are produced by zona fasciculata cells in adrenal. In the ovary, corpus luteum, where the granulosa cells are located, secretes progesterone. In addition, the testicular Leydig cells are able to produce testosterone. Since PAP7 is highly expressed in major steroidogenic tissues and is more concentrated in these steroid producing cells, PAP7 may be involved in steroid biosynthesis or the regulation of steroid formation by changing the formation or the conformation of the PBR complex. Mouse C6 glioma cells, MA-10 Leydig cells and Y1 adrenal cortical cells are popular cell models selected for studying steroid biosynthesis. PAP7 expression is proportionally correlated to the PBR expression in these cell lines. Additionally, both PBR and PAP7 expression level in these cell lines are parallel with their steroidogenic capability, which also suggests that PAP7 may be involved in steroid biosynthesis through a PBR pathway. A small PAP7 transcript was expressed only in testis, a phenomenon observed for other genes expressed in testis (Zhang, F. P. et al., 1997, *Endocrinology* 138, 2481-2490; Mauduit, C. et al., 1999, *J. Biol. Chem.* 274, 770-775). The immunostaining in the testis other than in Leydig cells could represent the expression of the smaller transcript.

The PAP7 protein is expressed in MA-10 cells and most of the staining is localized in the cytoplasm. The study of PAP7 subcellular distribution is ongoing,

the results of which may provide more detailed information about the interaction between PBR and PAP7. Since PBR knockout mice die in uterus indicates an essential role for PBR essential in mouse embryonic development. Interestingly, PAP7 mRNA is highly expressed during mouse early embryonic development. This result may suggest that PAP7, associated with PBR, could play an important role during early mouse development. This further implies that PBR may have new functions beyond steroidogenesis. Overexpression of PAP7 fragment including its PBR binding domain significantly inhibited the progesterone formation stimulated by saturating concentrations of hCG (50 ng/ml) in MA-10 cells. According to previous studies, progesterone production of these cells represented the index of steroid biosynthesis (Freeman, D. A., 1987, *Endocrinology* 120, 124-132; Garnier, M. et al., 1994, *J. Biol. Chem.* 269, 22105-22112). Based on the inhibitory manner, we assume that the overexpressed PAP7 fragment might act as a competitor of the native PAP7 in MA-10 cells and competitively bind to PBR. We believe that the tranfected PAP7 fragment having only the PBR binding domain and is not fully functional as a native PAP7, however, it competitively prevents PBR from interacting with the endogenous PAP7 and thereby blocks the normal function of PBR.

In conclusion, the results presented herein suggest that the identified PAP7 is involved in the regulation of the PBR function, serving as an endogenous ligand or allosteric modulator of the receptor.

Considering the findings tha i) PBR is a channel/transporter of cholesterol, ii) PBR is the target of environmental antisteroidogenic hazards, and (iii) PBR is involved in breast cancer aggression and

tumor cell proliferation (Hardwick, M. et al., 1999, *Cancer Res.* 59, 831-842), we believe that the identification and characterization of PAP7 will greatly contribute to the understanding of the role of
5 PBR in steroidogenesis and even in more general areas.

10

15

20

25

30

35

What is claimed is:

1. An isolated PBR-associated protein (PAP) DNA fragment or any portion thereof.
2. An isolated and purified DNA fragment which
5 encodes a PBR-associated protein.
3. An isolated and purified DNA fragment which encodes a peptide of PBR-associated protein, said DNA fragment comprising a sequence specified in any of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, and
10 SEQ ID NO:5, or polynucleotide fragment of said sequence comprising at least 30 nucleotides.
4. An isolated and purified DNA fragment which encodes a peptide of PBR-associated protein, said DNA fragment comprising a sequence specified in Genbank
15 Accession no. AF022770, or GenBank Accession no. AF020338, or a polynucleotide fragment of said sequence comprising at least 30 nucleotides.
5. An isolated and purified PAP7 DNA fragment according to claim 2 which encodes 463 amino acids of
20 PAP7 or a natural variant or synthetic variant thereof encoding PAP7, or a peptide fragment thereof comprising at least 10 amino acids.
6. An isolated and purified PAP8 DNA fragment according to claim 2 which encodes 190 amino acids of
25 PAP8 or a natural variant or synthetic variant thereof encoding PAP8, or a peptide fragment thereof comprising at least 10 amino acids.
7. An isolated and purified PAP15 DNA fragment according to claim 2 which encodes 164 amino acids of
30 PAP15 or a natural variant or synthetic variant thereof encoding PAP15, or a peptide fragment thereof comprising at least 10 amino acids.
8. An isolated and purified PAP20 DNA fragment according to claim 2 which encodes 196 amino acids of
35 PAP20 or a natural variant or synthetic variant

thereof encoding PAP20, or a peptide fragment thereof comprising at least 10 amino acids.

9. A recombinant DNA construct comprising:

(i) a vector, and

5 (ii) the PAP DNA fragment of claim 1.

10. A recombinant DNA construct comprising:

(i) a vector, and

(ii) the PAP DNA fragment of claim 3.

11. A recombinant DNA construct according to
10 claim 10, wherein said vector is an expression vector.

12. The recombinant DNA construct according to
claim 10, wherein said vector is a prokaryotic vector.

13. The recombinant DNA construct according to
claim 10, wherein said vector is a eukaryotic vector.

15 14. A host cell transformed with a recombinant
DNA construct according to claim 10.

15. A host cell according to claim 14, wherein
said cell is prokaryotic.

20 16. A host cell according to claim 14, wherein
said cell is eukaryotic.

17. A method for producing PAP peptide which
comprises culturing the cells according to either
claim 15 or 16, under conditions such that said DNA
fragment is expressed and said PAP peptide is thereby
25 produced.

18. An isolated recombinant PAP produced by the
method of claim 17.

19. A PAP7 polypeptide comprising the amino acid
sequence specified in SEQ ID NO:7 or a portion thereof
30 of at least 5 amino acids.

20. A PAP8 polypeptide comprising the amino acid
sequence specified in SEQ ID NO:8 or a portion thereof
of at least 5 amino acids.

21. A PAP15 polypeptide comprising the amino acid sequence specified in SEQ ID NO:9 or a portion thereof of at least 5 amino acids.

22. A PAP20 polypeptide comprising the amino acid
5 sequence specified in SEQ ID NO:10 or a portion thereof of at least 5 amino acids.

23. A method for detecting a PAP in a sample chosen from the group consisting of: PAP7 , PAP8, PAP15, PAP20, said method comprising

10 (i) contacting said sample with antibodies which recognize said PAP; and

(ii) detecting the presence or absence of a complex formed between PAP and antibodies specific therefor.

15 24. A method for detecting a PBR-associated protein, said method comprising the two hybrid assay.

25. An antibody to a peptide having the amino acid sequence specified in SEQ ID NO:6, 7, 8 and 9, or any portion thereof.

20 26. A PAP7 antibody to a peptide comprising the amino acid sequence specified in SEQ ID NO:11.

27. A method for detecting agents or drugs which reduce or eliminate PAP activity, said method comprising:

25 (i) delivering a recombinant DNA construct according to claim 10 into a cell such that PAP is produced in said cell;

(ii) adding at least one drug or agent to said cell alone or in combination; and,

30 (iii) detecting PAP activity in said cell in the presence of said agent or drug and comparing it to a control which did not receive said drug or agent wherein a decrease in PAP activity as compared to control indicates an drug or agent which reduces or
35 eliminates PAP activity.

28. A method for detecting agents or drugs which promote PAP activity, said method comprising:

- (i) delivering a recombinant DNA construct according to claim 10 into a cell such that PAP is
5 produced in said cell;
- (ii) adding at least one drug or agent to said cell alone or in combination; and,
- (iii) detecting whether or not said drug or agent stimulates PAP activity by measuring PAP activity in
10 said cell and comparing it to a control which did not receive said drug or agent wherein an increase in the activity of said PAP in said cell as compared to control indicates a stimulatory drug or agent.

29. An agent or drug capable of inhibiting PAP
15 activity.

30. An agent or drug capable of promoting PAP activity.

31. A therapeutic compound comprising said agent or drug according to claim 29 for use in a disease
20 wherein a decrease or elimination of PAP activity is beneficial.

32. A therapeutic compound comprising said agent or drug according to claim 30 for use in a disease wherein an increase o PAP activity is beneficial.

25 33. A method for detecting at least one PAP selected from the group consisting of PAP7, PAP8, PAP15, and PAP20 in a sample using the polymerase chain reaction.

34. A diagnostic kit for detecting RNA/cDNA of at
30 least one PAP chosen from the group consisting of PAP7, PAP8, PAP15 and PAP20, in a sample comprising primers or oligonucleotides specific for said PAP RNA or cDNA suitable for hybridization to PAP RNA or cDNA and/or amplification of PAP sequences and suitable
35 ancillary reagents.

35. A method for increasing a PAP selected from the group consisting of PAP3, PAP7, PAP8, PAP15 and PAP20 in a cell by introducing into said cell a PAP nucleic acid encoding said PAP such that said nucleic acid is expressed and PAP is produced in said cell.

36. A therapeutic method for the treatment or amelioration of diseases resulting from an increase in cell proliferation, said method comprising providing to an individual in need of such treatment an effective amount of an agent or drug which reduces or eliminates PAP expression or function in a pharmaceutically acceptable diluent.


37. The method of claim 36 wherein said disease is cancer.

38. A therapeutic method for the treatment or amelioration of conditions resulting from abnormal cholesterol level, said method comprising providing to an individual in need of such treatment an effective amount of an agent or drug which reduces or eliminates PAP expression or function in a pharmaceutically acceptable diluent.

39. The method of claim 38 wherein said condition is selected from the group consisting of: cancer, neurodegenerative disorders, developmental disorders, stress, and stroke.

40. A method for modulating PBR activity, function or targeting in a cell, said method comprising increasing or decreasing level of PAP in said cell.

A



B




FIGURE 1/8

A

	1	2	3	4	5
A	brain	eye	liver	lung	kidney
B	heart	skeletal muscle	small intest.		
C	pancreas	thyroid	thymus	submax. gland	spleen
D	testis	ovary	prostate	epidi-dymus	uterus
E	embryo 7 days	embryo 11 days	embryo 15 days	embryo 17 days	


B

FIGURE 2/8



A**B**

FIGURE 3/8



A**B**

FIGURE 4/8




FIGURE 5/8



A**B**

FIGURE 6/8

7 / 8





FIGURE 7/8

8 / 8





FIGURE 8/8

DAD20

SEQ ID NO:1

PAP3 nucleic acid

GAATTCGCGGCCGCGTCGACCACCGCTGCGCCCTCCTGCGAGGCCGGCTGAACGAGGAAATA
ATTGCTAATAAGGCCTCTGTAGCCATGGCTACTTCTGACGTGAAACCAAAATCAATAAGTCGTG
CCAAGAAATGGTCAGAGGAAATAGAAAATCTGTACAGATTTCAACAAGCAGGATATCGGGATG
AAATTGAATATAAACAAAGTGAACAAGTTGCCATGGTCGACCGATGGCCAGAGACAGGGTACG
TGAAGAACTTCAGCGGAGGGACAATACTTTCTTCTACTACAACAAAGAGAGGGAGTGCGA
GGACAAGGAGGTCCACAAAGTGAAGGTTTACGTCTACTGACCTTTTCTTTCTTCGGCTTG
GCAATGCTCCTTTAAGAATTGGTTGTTTACATTCTTCCATCGTGTAATGTCATTTTACAA
AACAAATTCACAATTCTGTCTTTAATTTCATGGTGTCTTACACAACATAAACACCCACCTTGA
AACCCAAAAA

SEQ ID NO:2

PAP7 sequence:

GAATTCGCGGCCGCGTCGACCTAAAGTTGAGTTGTTCACTGTAGTGACCCGTGTGAAGGTAGTT
TTATTTTTAAATCAACTTTTCATTGTGCAAACTAGTAAAAGATGGCAAAGCCTTTTCATCCAACTTA
TGAAGAAAAAAGTGAAGTTTCGTGGCACTGCATAAGCAAGTTCTTTTGGGCCCATAAACCCAGA
CACGTCCCCTGAGGTTGGATTCTTTGATGTGTTGGGGAATGATAGGAGGAGAGAATGGGCAGCT
CTGGGAAACATGTCCAAGGAGGATGCCATGGTAGAGTTTGTGAAGCTTCTAAATAAGTGTGTC
CTCTCCTCTCGGCATATGTTGCGTCCCACAGAATAGAGAAGGAAGAAGAAGAGAAAAAGAAGAA
AGGCGGAGGAGGAGCGAAGGCAGCGTGAAGAGGAAGAACGAGAGCGGCTGCAAAAGGAAGAA
GAGAAGCGGAAGCGAGAGGAGGAAGACCGGCTGAGACGGGAGGAGGAAGAGAGGCGGCGGAT
AGAGGAAGAGAGGCTTCGGCTGGAACAGCAAAAGCAGCAGATAATGGCAGCTTTAAACTCGCA
GACTGCCGTGCAATTCCAGCAGTATGCAGCCCAGCAGTATCCAGGGAACACGAACAACAGCA
GATTCTCATCCGCCAGCTGCAGGAGCAGCACTATCAGCAGTATAAACACCCAGGCAGAGCAAAC
CCAACCTGCACAACAACAGGCAGCATTACAGAAACAGCAAGAAGTAGTGATGGCTGGGGCAT
CATTGCCTGCATCATCAAAGGTGAACACAGCTGGAGCAAGTGATACACTGTCACTTAATGGAC
AGGCCAAAACCCACACTGAAAATTCCGAAAAAGTCCTTGAGCCAGAAGCTGCAGAAGAAGCC
TTGGAAAAATGGACCAAAAGACTCTCTTCCAGTGATTGCAGCTCCATCCATGTGGACAAGACCA
CAAATCAAAGACTTTAAAGAGAAGATTCCGGCAGGATGCAGATTCTGTGATTACAGTACGTCCA
GGAGAAGTCGTACCGTCCGAGTCCCGACTCATGAGGAAGGATCATACCTATTTTGGGAATTTG
CCACAGACAGTTATGACATTGGGTTTGGGGTTTATTTTGAATGGACAGACTCTCCAAATGCTGC
TGTCAGTGTGCATGTCACTGAGTCCAGTGACGAGGAGGAGGAGGAGGAAGAAAATGTCACTTG
TGAAGAAAAAGCAAAAAAAGAACGCCAACAAAGCCTCTGCTGGATGAGATTGTACCTGTGTACCG
GCGGACTGTACGAGGAAGTATATGCAGGCAGCCACCAGTATCCAGGGAGGGAGTCTATC
TCCTCAAGTTTGATAATTCTTACTCTCTGTGGAGGTCCTCAAGTCCGTCTACTACAGAGTCTA
TTATACTAGATAGAGCTGCTGTTCCAAGGTCCGGAGTCCAGGGTTGAGCACAACATGACGT
TTAATTTCTCTT

SEQ ID NO:3

PAP8 nucleic acid

GAATTCGCGGCCGCGTCGACGCTGGACACAAGCGTGGAGCGAAGAGCCCTTGGAGAGATTTCAG
AATGTAGTGGAAGGCTCTTCAACATCACAGGGCACCTGGCAGTCTTCAGAGTCTTCACAGTCA
AACCTGGGGGAGCAGACGCAGAGCGGACCCACGGAGGAAGGTGTCAGCGTCCGGGAGAGGCA
TAACCGAATGGAACGAGATAGGAGGCGCAGAATCCGCATTTGCTGTGATGAGCTGAATCTTTTA
GTTCCATTCTGCAATGCGGAGACAGATAAAGCAACAACCCTTCAGTGGACCACAGCATTCTCTG
AAGTACATTTCAGGAAAGACATGGGGACTCTCTTAAAAAAGGAATTTGAGAGCGTGTGTTGCGGTA
AAACTGGCAGAAGGCTAAAGCTGACTAGACCCGAATCCCTGGTGACCTGCCCTGCACAGGGCA
GCCTGCAGAGCAGCCCTGCCATGGAGATCAAGTGACTGGACTGACCCAGGACCTGGGAGAGA

ACCGCCGTTCTCTGCGGCATCATGCACATGCCTGCCATCCCCGGAATTCAGCTCTGAATCCT
CTC

SEQ ID NO:4

PAP15 nucleic acid

GAATTCGCGGCCGCGTCGACCTCGGGGGTAACCCTTGGGATCATTATGATGTCACCTCTTCAAT
CTCTTGAGACCAGTGATTTATGAACTTGACATTGGTGCCTGGTGTGGGCTTTGGAAAGCAGAG
GCCACCTTTGTCTGTAGAGGATACTGAGCGGCTGGATGGCAGGAATCCAAAGAGAAGCCAGCC
TCTCGTAGTCGCCCTGGGACAGTGGAAGAGAGGGAGGACCGGCAAAGGGGAATCTGTCTTTCT
CCTAGGCCCGAGCATGTCCCCTGTGGGACATGCTCTGTGACAGCTGAGCCTGCCCAGCCTG
CCTTTCTGAAGTTGGGTGTCTCCTGCCCACAACCAAGCCAGCAATCGGTCTGTTTTCCGAC
AACCTCAGAGCCAGACCTCACAAGCCTATTTTGGTGGTTCCCAAATTTCTCTCAGATCTC
CATGTCTATCCCTCCACTCCCTCCAAAAGAGAAAGAAAAGAATTGAGAAAGAA

SEQ ID NO:5

PAP20 nucleic acid

GAATTCGCGGCCGCGTCGACCTCAGAAGAGGAAAAAGAGGTGCAGAAGTGCTGGCGGCACAAA
TTGTACAGAAAACCAGACTAGAGAGAAAAAAACAAGAAGCGTCTGTATCTAAAGATGCTCCAG
TGCCTACAAATACTAAAAGGGCAAAGAAACAAGAGAAGTCTCCAGGTAGAATTGCCTCACAGT
CTAAGCCACCCATGAAAAAGTCTCCACAAAAACGGAAGGTAAATGTAGCAAGAGGCCGTCGG
AATACCAGAAAAGCAGTCCAACCTGCCGAAAAAGAAATTGCTTTACATCTTCAATCAGAAATT
TCATCAGATGGCCAAAAAGATGGACTTAACCTAAGTACATCTCAACAAGAAAGTATTTCAATG
ATTCTAAAGGTCCTCCTGAAAACTCAGTTATCAGCTGTGACTCCCAGGCCCTAAATATGTTAG
CTGATCTGGCATTAAAGTTCTGCTGCTGCTTCTATACCATCTTGTAAGCCCAGGAACCTTCCCTG
CGTCTCTGATTTGCCACGAAACAATGTCTTACTCACTAAAGAAAATCCATTGCTTGGTGCCT
CTGACCATGAATATCATAAGGG

SEQ ID NO:6

PAP3 amino acid sequence

Met Ala Thr Ser Asp Val Lys Pro Lys Ser Ile Ser Arg Ala Lys
Lys Trp Ser Glu Glu Ile Glu Asn Leu Tyr Arg Phe Gln Gln Ala
Gly Tyr Arg Asp Glu Ile Glu Tyr Lys Gln Val Lys Gln Val Ala
Met Val Asp Arg Trp Pro Glu Thr Gly Tyr Val Lys Lys Leu Gln
Arg Arg Asp Asn Thr Phe Phe Tyr Tyr Asn Lys Glu Arg Glu Cys
Glu Asp Lys Glu Val His Lys Val Lys Val Tyr Val Tyr

SEQ ID NO:7

PAP7 amino acid sequence

Arg Pro Arg Arg Pro Lys Val Glu Leu Phe Thr Val Val Thr Arg
Val Lys Val Val Leu Phe Leu Asn Gln Leu Ser Leu Cys Lys Leu
Val Lys Asp Gly Lys Ala Phe His Pro Thr Tyr Glu Glu Lys Leu
Lys Phe Val Ala Leu His Lys Gln Val Leu Leu Gly Pro Tyr Asn
Pro Asp Thr Ser Pro Glu Val Gly Phe Phe Asp Val Leu Gly Asn
Asp Arg Arg Arg Glu Trp Ala Ala Leu Gly Asn Met Ser Lys Glu
Asp Ala Met Val Glu Phe Val Lys Leu Leu Asn Lys Cys Cys Pro

Leu Leu Ser Ala Tyr Val Ala Ser His Arg Ile Glu Lys Glu Glu
Glu Glu Lys Arg Arg Lys Ala Glu Glu Arg Arg Gln Arg Glu
Glu Glu Glu Arg Glu Arg Leu Gln Lys Glu Glu Lys Arg Lys
Arg Glu Glu Glu Asp Arg Leu Arg Arg Glu Glu Glu Arg Arg
Arg Ile Glu Glu Glu Arg Leu Arg Leu Glu Gln Gln Lys Gln Gln
Ile Met Ala Ala Leu Asn Ser Gln Thr Ala Val Gln Phe Gln Gln
Tyr Ala Ala Gln Gln Tyr Pro Gly Asn Tyr Glu Gln Gln Gln Ile
Leu Ile Arg Gln Leu Gln Glu Gln His Tyr Gln Gln Tyr Lys His
Gln Ala Glu Gln Thr Gln Pro Ala Gln Gln Gln Ala Ala Leu Gln
Lys Gln Gln Glu Val Val Met Ala Gly Ala Ser Leu Pro Ala Ser
Ser Lys Val Asn Thr Ala Gly Ala Ser Asp Thr Leu Ser Val Asn
Gly Gln Ala Lys Thr His Thr Glu Asn Ser Glu Lys Val Leu Glu
Pro Glu Ala Ala Glu Glu Ala Leu Glu Asn Gly Pro Lys Asp Ser
Leu Pro Val Ile Ala Ala Pro Ser Met Trp Thr Arg Pro Gln Ile
Lys Asp Phe Lys Glu Lys Ile Arg Gln Asp Ala Asp Ser Val Ile
Thr Val Arg Arg Gly Glu Val Val Thr Val Arg Val Pro Thr His
Glu Glu Gly Ser Tyr Leu Phe Trp Glu Phe Ala Thr Asp Ser Tyr
Asp Ile Gly Phe Gly Val Tyr Phe Glu Trp Thr Asp Ser Pro Asn
Ala Ala Val Ser Val His Val Ser Glu Ser Ser Asp Glu Glu Glu
Glu Glu Glu Glu Asn Val Thr Cys Glu Glu Lys Ala Lys Lys Asn
Ala Asn Lys Pro Leu Leu Asp Glu Ile Val Pro Val Tyr Arg Arg
Asp Cys His Glu Glu Val Tyr Ala Gly Ser His Gln Tyr Pro Gly
Arg Gly Val Tyr Leu Leu Lys Phe Asp Asn Ser Tyr Ser Leu Trp
Arg Ser Lys Ser Val Tyr Tyr Arg Val Tyr Tyr Thr Arg

SEQ ID NO:8

PAP8 amino acid sequence

Glu Phe Ala Ala Ala Ser Thr Leu Asp Thr Ser Val Glu Arg Arg
Ala Leu Gly Glu Ile Gln Asn Val Gly Glu Gly Ser Ser Thr Ser
Gln Gly Thr Trp Gln Ser Ser Glu Ser Ser Gln Ser Asn Leu Gly
Glu Gln Thr Gln Ser Gly Pro Gln Gly Gly Arg Cys Gln Arg Arg
Glu Arg His Asn Arg Met Glu Arg Asp Arg Arg Arg Arg Ile Arg
Ile Cys Cys Asp Glu Leu Asn Leu Leu Val Pro Phe Cys Asn Ala
Glu Thr Asp Lys Ala Thr Thr Leu Gln Trp Thr Thr Ala Phe Leu
Lys Tyr Ile Gln Glu Arg His Gly Asp Ser Leu Lys Lys Glu Phe
Glu Ser Val Phe Cys Gly Lys Thr Gly Arg Arg Leu Lys Leu Thr
Arg Pro Glu Ser Leu Val Thr Cys Pro Ala Gln Gly Ser Leu Gln
Ser Ser Pro Ala Met Glu Ile Lys

SEQ ID NO:9

PAP15 amino acid sequence

Ala Ala Gly Trp Gln Glu Ser Lys Glu Lys Pro Ala Ser Arg Ser
Arg Pro Gly Thr Val Glu Glu Arg Glu Asp Arg Gln Arg Gly Ile
Cys Leu Ser Pro Arg Pro Glu His Val Pro Cys Gly Thr Cys Ser
Val Thr Ala Glu Pro Ala Gln Pro Ala Phe Leu Lys Leu Gly Val
Ser Cys Pro Gln Pro Ser Gln Gln Ser Val Cys Phe Pro Thr Thr
Ser Glu Pro Asp Leu Thr Ser Leu Phe Trp Trp Phe Pro Lys Phe
Leu Ser Asp Leu His Val Tyr Pro Ser Thr Pro Ser Lys Arg Glu
Arg Lys Glu Leu Arg Lys Lys

SEQ ID NO:10

PAP20 amino acid sequence

Asn Ser Arg Pro Arg Arg Pro Gln Lys Arg Lys Arg Gly Ala Glu
Val Leu Ala Ala Gln Ile Val Gln Lys Thr Arg Leu Glu Arg Lys
Lys Gln Glu Ala Ser Val Ser Lys Asp Ala Pro Val Pro Thr Asn
Thr Lys Arg Ala Lys Lys Gln Glu Lys Ser Pro Gly Arg Ile Ala
Ser Gln Ser Lys Pro Pro Met Lys Lys Ser Pro Gln Lys Arg Lys
Val Asn Val Ala Arg Gly Arg Arg Asn Thr Arg Lys Gln Leu Gln
Pro Ala Glu Lys Glu Ile Ala Leu His Leu Gln Ser Glu Ile Ser
Ser Asp Gly Gln Lys Asp Gly Leu Asn Leu Ser Thr Ser Gln Gln
Glu Ser Ile Ser Met Ile Pro Lys Gly Pro Pro Glu Asn Ser Val
Ile Ser Cys Asp Ser Gln Ala Leu Asn Met Leu Ala Asp Leu Ala
Leu Ser Ser Ala Ala Ala Ser Ile Pro Ser Cys Lys Pro Arg Asn
Leu Pro Cys Val Ser Asp Leu Pro Arg Asn Asn Val Leu Leu Thr
Lys Glu Asn Pro Leu Leu Gly Ala Ser Asp His Glu Tyr His Lys
Gly

SEQ ID NO:11

Ser Ser Asp Glu Glu Glu Glu Glu Glu Glu Asn Val Thr Cys Glu
Glu Lys Ala Lys Lys Asn Ala Asn Lys Pro



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07K 14/00	A2	(11) International Publication Number: WO 00/09549 (43) International Publication Date: 24 February 2000 (24.02.00)
(21) International Application Number: PCT/US99/18507 (22) International Filing Date: 11 August 1999 (11.08.99) (30) Priority Data: 60/096,048 11 August 1998 (11.08.98) US <i>11 Feb 01 / 30 MOS</i> (71) Applicant (for all designated States except US): GEORGETOWN UNIVERSITY MEDICAL CENTER [US/US], Office of the Dean of Research and Graduate Education, Building D, suite 177, Washington, DC 20007 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): PAPAPOPOULOS, Vasilios [US/US]; 15417 Peach Leaf Drive, North Potomac, MD 20878 (US). LI, Hua [CN/US]; Apartment 102, 1319 Fort Myer Drive, Arlington, VA (US). (74) Agent: PRATT, Sana, A.; Pratt & Associates, Inc., 10821 Hillbrooke Lane, Potomac, MD 20854 (US).		(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: PERIPHERAL-TYPE BENZODIAZEPINE RECEPTOR ASSOCIATED PROTEINS, CLONING, EXPRESSION AND METHODS OF USE		
(57) Abstract <p>The present invention relates to nucleic acids encoding PBR-associated proteins (PAPs) and methods for use in producing PAPs and methods of using PAPs. In this study we have identified proteins (PAPs) that interact with PBR protein by a yeast two-hybrid system. We used PBR as bait to screen a mouse testis cDNA library. Five clones were isolated based on their ability to interact with PBR. These proteins may be involved in the regulation of the function of PBR, serving as an endogenous ligand or allosteric modulator of the receptor.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE OF THE INVENTION

Peripheral-type Benzodiazepine Receptor Associated
Proteins, Cloning, Expression and Methods of Use

5

Field of the Invention

The present invention relates to nucleic acid molecules encoding peripheral-type benzodiazepine receptor (PBR) associated proteins (PAPs), including
10 mutants, variants, fragments and derivatives thereof, and to vectors and host cells comprising such nucleic acid molecules; methods of using PAPs; method for screening for inhibitors and activators of PAPs or PBR; and kits comprising the compositions or
15 polypeptides of the invention.

Background of the Invention

The peripheral-type benzodiazepine receptor (PBR) was originally discovered because it binds the
20 benzodiazepine diazepam with relatively high affinity (Papadopoulos, V. 1993, *Endocr. Rev.* 14:222-240). Benzodiazepines are among the most highly prescribed drugs due to their pharmacological actions in relieving anxiety mediated through modulating the
25 activity of γ -aminobutyric acid receptors in the central nervous system (Costa, E. and Guidotti, A. 1979, *Annu. Rev. Pharmacol. Toxicol.* 19:531-545). PBR is another class of binding sites for benzodiazepines distinct from the aforementioned neurotransmitter
30 receptors. Further studies demonstrated that in addition to benzodiazepines, PBR binds other classes of organic compounds with high affinity (Papadopoulos, 1993, *supra*). PBR, although present in all tissues examined, was found to be particularly high in steroid

producing tissues, where it was primarily localized in the outer mitochondrial membrane (OMM) (Anholt, R.R.H. et al. 1986, *J. Biol. Chem.* 261:576-583). An 18 kDa isoquinoline-binding protein was identified as PBR, cloned and expressed (Papadopoulos, V. 1998, *Proc Soc. Exp. Biol. Med.* 217:130-142). It was then demonstrated that PBR is a functional component of the steroidogenic machinery (Papadopoulos, 1998, *supra*; Papadopoulos V. et al. 1990, *J. Biol. Chem.* 265:3772-3779) mediating cholesterol delivery from the outer to the inner mitochondrial membrane (Krueger, K. E. and Papadopoulos, V. 1990, *J. Biol. Chem.* 265:15015-15022). Further studies demonstrated that pharmacologically induced reduction of adrenal PBR levels *in vivo* resulted in decreased circulating glucocorticoid levels (Papadopoulos, V. 1998, *supra*). In addition, targeted disruption of the PBR gene in Leydig cells resulted in the arrest of cholesterol transport into mitochondria and steroid formation; transfection of the mutant cells with a PBR cDNA rescued steroidogenesis (Papadopoulos, V. et al. 1997, *J. Biol. Chem.* 272:32129-32135).

PBR is extremely abundant in steroidogenic cells and found primarily on outer mitochondrial membranes (Anholt, R. et al. 1986, *J. Biol. Chem.* 261:576-583). PBR is thought to be associated with a multimeric complex composed of the 18-kDa isoquinoline-binding protein and the 34-kDa pore-forming voltage-dependent anion channel protein, preferentially located on the outer/inner mitochondrial membrane contact sites (McEnery, M. W. et al. *Proc. Natl. Acad. Sci. U.S.A.* 89:3170-3174; Garnier, M. et al. 1994, *Mol. Pharmacol.* 45:201-211; Papadopoulos, V. et al. 1994, *Mol. Cel. Endocr.* 104:R5-R9). Drug ligands of PBR, upon binding to the receptor, simulate steroid synthesis in

steroidogenic cells *in vitro* (Papadopoulos, V. et al. 1990, *J. Biol. Chem.* 265:3772-3779; Ritta, M. N. et al. 1989, *Neuroendocrinology* 49: 262-266; Barnea, E. R. et al. 1989, *Mol. Cell. Endocr.* 64:155-159;

5 Amsterdam, A. and Suh, B. S. 1991, *Endocrinology* 128:503-510; Yanagibashi, K. et al. 1989, *J. Biochem. (Tokyo)* 106: 1026-1029). Likewise, *in vivo* studies showed that high affinity PBR ligands increase steroid plasma levels in hypophysectomized rats (Amri, H. et

10 al. 1996, *Endocrinology* 137:5707-5718). Further *in vitro* studies on isolated mitochondria provided evidence that PBR ligands, drug ligands, or the endogenous PBR ligand, the polypeptide diazepam-binding inhibitor (BDI) (Papadopoulos, V. et al. 1997,

15 *Steroids* 62:21-28), stimulate pregnenolone formation by increasing the rate of cholesterol transfer from the outer to the inner mitochondrial membrane (Krueger, K. E. and Papadopoulos, V. 1990, *J. Biol. Chem.* 265:15015-15022; Yanagibashi, K. et al. 1988,

20 *Endocrinology* 123: 2075-2082; Besman, M. J. et al. 1989, *Proc. Natl. Acad. Sci. U.S.A.* 86: 4897-4901; Papadopoulos, V. et al. 1991, *Endocrinology* 129: 1481-1488).

Based on the amino acid sequence of the 18-kDa

25 PBR, a three dimensional model was developed (Papadopoulos, V. 1996, In: The Leydig Cell. Payne, A. H. et al. (eds) Cache River Press, IL, pp 596-628). This model was shown to accomodate a cholesterol molecule and function as a channel, supporting the

30 role of PBR in cholesterol transport. Recently we demonstrated the role of PBR in steroidogenesis by generating PBR negative cells by homologous recombination (Papadopoulos, V. et al. 1997, *J. Biol. Chem.* 272:32129-32135) that failed to produce

35 steroids. However, addition of the hydrosoluble

analogue of cholesterol, 22R-hydroxycholesterol, recovered steroid production by these cells, indicating that the cholesterol transport mechanism was impaired. Further cholesterol transport experiments in bacteria expressing the 18-kDa PBR protein provided definitive evidence for a function as a cholesterol channel/transporter (Li and Papadopoulos, 1998, *Endocrinology* 139, 4991-4997).

Studies in a number of tumors such as rat brain containing glioma tumors (Richfield, E. K. et al. 1988, *Neurology* 38:1255-1262), colonic adenocarcinoma and ovarian carcinoma (Katz, Y. et al. 1988, *Eur. J. Pharmacol.* 148: 483-484 and Katz, Y. et al. 1990, *Clinical Sci.* 78:155-158) have shown an abundance of peripheral-type benzodiazepine receptors (PBR) compared to normal tissue. All documents cited herein *infra* and *supra* are hereby incorporated in their entirety by reference thereto. Moreover, a 12-fold increase in PBR density relative to normal parenchyma, was found in human brain glioma or astrocytoma (Cornu, P. et al. 1992, *Acta Neurochir.* 119:146-152). The authors suggested that PBR densities may reflect the proliferative activity of the receptor in these tissues. Recently, the involvement of PBR in cell proliferation was further shown (Neary, J. T. et al. 1995, *Brain Research* 675:27-30; Miettinen, H. et al. 1995, *Cancer Research* 55:2691-2695), and its expression in human astrocytic tumors was found to be associated with tumor malignancy and proliferative index (Miettinen, H. et al. *supra*; Alho, H. 1994, *Cell Growth Different.* 5:1005-1014). Characterization of PBR in human breast cancer biopsies, led to the discovery that the invasive and metastatic ability of human breast tumor cells is proportional to the level of PBR expressed, and correlates with the subcellular

localization of PBR in these cells in that PBR is found primarily in the nucleus in aggressive tumor cells whereas PBR is found primarily in the cytoplasm of invasive but non-aggressive cells. These changes in PBR expression can be used as a tool for detection, diagnosis, prevention and treatment in breast cancer patients, in particular, and in aggressive solid tumors in general.

Since both PBR and its endogenous ligand, the polypeptide diazepam binding inhibitor, are constitutively expressed in steroidogenic cells, the regulation of PBR function by hormones may be due to its association with other proteins. This interaction may result in the initiation of steroid biosynthesis. Therefore, there is a need to identify proteins which associate with PBR and may modulate PBR function.

Summary of the Invention

The present invention meets the need described above. We have identified PBR associated proteins (PAPs) that interact with PBR using the two-hybrid system. We used PBR as a bait to screen a mouse testis cDNA library. Five clones were isolated by their ability to interact with PBR: PAP3, PAP7, PAP8, PAP15, and PAP20. Among the nucleotide sequences identified, PAP3 was identical to the previously isolated meg1 protein (Don, J. and Wolgemuth, D. J., 1992, *Cell Growth Differ.* 3, 495; Ever, L. et al., 1999, *Cell Growth Differ.* 10, 19-26). PAP7, PAP8, PAP15, and PAP20 are novel sequences according to a search in the Genebank database which did not find a match for these sequences. PAP7 and PAP17 are different clones of the same novel protein product. All PAPs have fatty acylation (myristoylation) sites and PKC

phosphorylation sites. In addition, PAP20 has a PKA phosphorylation site. The distribution and function of the PAPs, as well as their functional relationship to PBR is under investigation.

5 So far, the distribution of PAP7 in major mouse tissues such as brain, testis, ovary, adrenal, kidney and muscle showed a profile similar to the broader expression pattern of PBR, with an expression level paralleling the steroidogenic ability of the tissue.
10 These data imply a role for these PAPs in the regulation of PBR function, serving as endogenous ligands or allosteric modulators of the receptor.

 Therefore, it is an object of the present invention to provide novel DNA fragments encoding PBR
15 associated proteins, PAP3 (SEQ ID NO:1), PAP7 (SEQ ID NO:2) and Genbank accession no. AF022770, PAP8 (SEQ ID NO:3), PAP15 (SEQ ID NO:4), and PAP20 (SEQ ID NO:5) GenBank accession no. AF020338. The DNA fragment is useful as a diagnostic agent for detection of nucleic
20 acid sequences encoding PBR-associated proteins, as an agent for preparation of the protein encoded by the DNA, for the preparation of sequences encoding PAPs, whether cDNA or genomic, and as therapeutic agents.

 It is another object of the invention to provide
25 an amino acid sequence for PAPs encoded by the DNA sequences described above.

 It is another object of the invention to provide a recombinant vector comprising a vector and the above described DNA fragments.

30 It is a further object of the present invention to provide a host cell transformed with the above-described recombinant DNA construct.

It is another object of the present invention to provide a method for producing PAPs which comprises culturing a host cell under conditions such that the above-described DNA fragment is expressed and a PAP is
5 thereby produced, and isolating the PAP for use as a reagent, for example for screening drugs and inhibitors of PBR or the PAP itself, for diagnosis, and for therapy.

It is a further object of the present invention
10 to provide an antibody to the above-described recombinant PAPs.

It is yet another object of the present invention to provide a method for detecting any of PAP3, PAP7, PAP8, PAP15, or PAP20 in a sample comprising:

- 15 (i) contacting a sample with antibodies which recognize any one of the PAPs mentioned above; and
(ii) detecting the presence or absence of a complex formed between the PAP and antibodies specific therefor.

20 It is a further object of the present invention to provide a diagnostic kit comprising an antibody against PAP and ancillary reagents suitable for use in detecting the presence of PAP in cells, tissue or serum from yeast, mammals, animals, birds, fish, and
25 plants.

It is yet another object of the present invention to provide a method for the detection of PAP from a sample using the polymerase chain reaction.

It is a further object of the present invention
30 to provide a diagnostic kit comprising primers or oligonucleotides specific for PAP RNA or cDNA suitable for hybridization to PAP RNA or cDNA and/or amplification of PAP sequences and ancillary reagents suitable for use in detecting PAP RNA/cDNA in
35 mammalian tissue.

It is yet another object of the present invention to provide a method for the detection of a PAP in a sample which comprises assaying for the presence or absence of PAP RNA or cDNA in a sample by
5 hybridization assays.

It is an object of the present invention to provide a method for the measurement of PBR in a sample. The method comprises measuring the presence of a PAP complexed with PBR.

10 It is yet another object of the present invention to provide a method for modulating the function or altering the targeting of PBR by increasing or decreasing interaction of PBR with a PAP. PBR functions which can be modulated include cholesterol
15 transport into the cell, steroid production, cell proliferation, and embryogenesis.

It is still another object of the present invention to provide a method for increasing or decreasing PBR function or expression in a cell by
20 providing into the cell a PAP such that PBR function is increased or decreased.

It is yet another object of the present invention to provide a method for increasing or reducing steroidogenesis by altering the level of PAP in said
25 cell.

It is still another object of the invention to provide a therapeutic method for the treatment or amelioration of diseases resulting from increased cell proliferation resulting from aberrant PBR function or
30 expression or localization, said method comprising providing to an individual in need of such treatment an effective amount of a PAP, in a pharmaceutically acceptable diluent, of a PAP such that the aberrant PBR expression, function or localization, is
35 corrected.

It is another object of the present invention to provide a therapeutic method for the treatment or amelioration of diseases resulting from decreased cell proliferation, said method comprising providing to an individual in need of such treatment an effective amount of PAP or an antibody against PAP or an agent which inhibits PAP expression or function in a pharmaceutically acceptable excipient.

It is another object of the present invention to provide a therapeutic method for the treatment or amelioration of diseases resulting from decreased or increased steroidogenesis, said method comprising providing to an individual in need of such treatment an effective amount of a PAP or an antibody against PAP or an agent which inhibits or activates PAP expression or function in a pharmaceutically acceptable diluent.

It is yet a further object of the present invention to provide a cDNA sequence encoding PAPs and vectors incorporating all or a fragment of said sequence, and cells, prokaryotic and eukaryotic, transformed or transfected with said vectors, for use in screening agents and drugs which inhibit expression or function of PAPs or PBR in such cells.

25

BRIEF DESCRIPTION OF THE DRAWINGS

These and other features, aspects, and advantages of the present invention will become better understood with reference to the following description and appended claims, and accompanying drawings where:

Figure 1. PAP7 protein expressed in MA10 mouse Leydig tumor cells. (A) Western blot. (B), Immunocytochemistry.

Figure 2. PAP7 mRNA Tissue distribution analysis by Dot blot analysis. (A), a Master blot

35

containing 100-500 ng of poly(A)+ RNA from mouse tissues were hybridized at high stringency with a ^{32}P -labeled PAP7 probe as described under "Materials and Methods." The autoradiogram was exposed overnight.

5 (B), densitometric analysis of PAP7 expression.

Figure 3. PAP7 tissue distribution analysis by Northern blot analysis. (A), Northern blot analysis was performed using 20 μg of total RNA/lane from different mouse tissues as indicated. The blot was
10 hybridized at high stringency with a ^{32}P -labeled PAP7 probe as described under "Materials and Methods". The autoradiogram was exposed overnight. (B), Densitometric analysis of PAP7 expression.

Figure 4. PBR tissue distribution analysis by
15 Northern blot analysis. (A), Northern blot analysis was performed using 20 μg of total RNA/lane from different mouse tissues as indicated. The blot was hybridized at high stringency with a ^{32}P -labeled PBR probe as described under "Materials and Methods". The
20 autoradiogram was exposed overnight. (B), For Northern blot analysis, blots were quantitated by densitometry.

Figure 5. Immunohistological staining of mouse tissues with anti-PAP7 antibody.

25 **Figure 6.** The effect of PAP7 on Steroid Biosynthesis. Progesterone formation in MA-10 stimulated by hCG (A), for different time course (B). The results shown represent the means + S.D. from 2 to 6 independent experiments.

30 **Figure 7.** PAP3 mRNA Tissue distribution analysis by Dot blot analysis. (A), a Master blot containing 100-500 ng of poly(A)+ RNA from mouse tissues were hybridized at high stringency with a ^{32}P -labeled PAP3 probe as described under "Experimental

Procedures." The autoradiogram was exposed overnight.
(B), densitometric analysis of PAP3 expression.

Figure 8. PAP20 mRNA Tissue distribution analysis by Dot blot analysis. (A), a Master blot
5 containing 100-500 ng of poly(A)+ RNA from mouse tissues were hybridized at high stringency with a ³²P-labeled PAP20 probe as described under "Experimental Procedures." The autoradiogram was exposed overnight.
(B), densitometric analysis of PAP20 expression.

10 **DETAILED DESCRIPTION**

The five PAPs described in this application were discovered using a two-hybrid assay. The two-hybrid assay is a yeast-based genetic assay used to detect protein-protein interactions *in vivo*. A positive
15 result obtained with the two-hybrid assay allows rapid identification of genes encoding proteins that interact with a target protein. In addition the two-hybrid assay is a sensitive method for detecting weak and transient interactions, which are probably the
20 norm in large native complexes. Most notably, because the two-hybrid assay is performed *in vivo*, the proteins involved are more likely to be in their native conformations.

The two-hybrid assay is based on the fact that
25 many eukaryotic transcriptional activators consist of two physically separable modular domains: one acts as the DNA-binding domain, while the other functions as the transcriptional activation domain. The DNA-binding domain localizes the transcription factor to
30 specific DNA sequences present in the upstream region of genes that are regulated by this factor, while the activation domain contacts other components of the transcription machinery required to initiate transcription. Both domains are required for normal

activation functioning, and normally the two domains are part of the same protein.

In our PAPs screening experiment, the MATCHMAKER Two-Hybrid System from CLONTECH was used. In the MATCHMAKER System, sequences encoding the two functional domains of the GAL4 transcriptional activator have been cloned into two different shuttle/expression vectors (pGBT9 and pGAD10). The pGBT9 hybrid cloning vector is used to generate a fusion of the GAL4 DNA-binding domain with PBR protein. The pGAD10 hybrid cloning vector is used to generate a fusion of the GAL4 activation domain with a collection of random proteins in a fusion mouse testis library (CLONTECH). Both hybrid proteins are targeted to the yeast nucleus by nuclear localization sequences that either are an intrinsic part of the GAL4 DNA-binding domain or have been added to the activation domain from a heterologous source. If PBR protein and an unknown protein or proteins interact with each other, the DNA-binding domain of GAL4 will be tethered to its transcriptional activation domain, and the proper function of the transcription of an appropriate reporter gene (lacZ or HIS3) containing upstream GAL4 binding sites is used to indicate interaction between the two proteins. This allows a positive selection for clones that are transformed by two interacting hybrid constructs and makes library screening more convenient and practical. After a positive clone has been identified, the gene corresponding to the interacting protein was sequenced using the sequencing primers provided in the kit.

In one embodiment, the present invention relates to a DNA or cDNA sequence encoding PBR associated proteins (PAPs). Five clones were isolated, PAP3, comprising 568 bp and identified in SEQ ID NO:1 (Don,

J. and Wolgemuth, D. J., 1992, *Cell Growth Differ.* 3, 495; Ever, L. et al., 1999, *Cell Growth Differ.* 10, 19-26) which encodes a peptide of 83 amino acids identified in SEQ ID NO:6; PAP7, comprising 577 bp
5 extending from 696 to 1164 of the sequence identified in SEQ ID NO:2, which encodes a polypeptide of 363 amino acids, identified in SEQ ID NO:7; PAP8, comprising 568 bp identified in SEQ ID NO:3, which encode a polypeptide of 190 amino acids, identified in
10 SEQ ID NO:8; PAP15 comprising 490 bp identified in SEQ ID NO:4, which encode a polypeptide of 164 amino acids, identified in SEQ ID NO:9; and PAP20 comprising 588 bp identified in SEQ ID NO:5, which encode a polypeptide of 196 amino acids, identified in SEQ ID
15 NO:10.

PAP3 has been identified as the previously isolated meg 1 protein.

PAP7 and PAP17 are different clones of the same novel protein product. Additional PAP7 sequence has
20 been obtained using the 5', 3'RACE system (CLONTECH) and the near full-length gene is identified in SEQ ID NO:2 including the stop codon and some untranslated sequence at the 3' end. The polypeptide encoded by the DNA sequence would have a calculated molecular
25 weight of about 50 kD. Using an PAP7 antibody produced from the initial isolated DNA fragment of 577 bp, a protein of about 52 kD is immunoprecipitated as shown in the Examples below. Analysis of the protein sequence indicates several consensus sequences and
30 important sites such as: two potential myristoylation sites at positions 262-267 and 271-276 of SEQ ID NO:7 and five PKC phosphorylation sites at 395-396, 113-115, 255-257, 280-282, 331-333, and 339-341 of SEQ ID NO:7, an Acyl-Co-A site at position 24-108 of SEQ ID
35 NO:7, a nuclear localization domain at position 150-

167 of SEQ ID NO:7, a troponin site at position 98-247 of SEQ ID NO:2 and an HSP90 domain at position 126-155 of SEQ ID NO:7. The distribution and expression of PAP7 were examined in major mouse tissues such as
5 brain, testis, ovary, adrenal, and kidney, as well as in tissue culture cell lines such as mouse C6 glioma cells, MA-10 Leydig cells, and Y1 adrenal cortical cells. The PAP7 expression pattern is similar to the broader expression profile of PBR in both tissues and
10 cell lines involved in steroid biosynthesis.

Additionally, both PBR and PAP7 expression level in the cell lines correlated with their steroidogenic biosynthesis ability, which suggests that PAP7 may be involved in steroid biosynthesis through PBR.

15 PAP 8, PAP15, and PAP20, are novel genes. The polypeptide encoded by PAP20 has two potential myristoylation sites, one PKC phosphorylation site and one PKA phosphorylation site. Protein myristoylation enables the protein to attach to the cellular membrane
20 and thus take part in cell signaling (Casey, P.J. 1995, *Science* 268, 221-225; Boutin, J. A. 1997, *Cell Signal* 9, 15-35). PAP20 is predominantly expressed in the testis. Interaction of PBR with PAP20 increased the affinity of ligand binding using PK11195 as a
25 ligand. Therefore, it is likely that PAP20 serves to increase or decrease PBR function by modulating PBR's affinity to its endogenous ligand, DBI. The tissue distribution of PAP3 and PAP20 is shown in Figure 7 and Figure 8, respectively.

30 Thus, one aspect of the invention provides an isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence selected from SEQ ID NO:1-9 encoding PAPs polypeptides. It is within the skill of a person with ordinary skill in
35 the art to use the sequences provided herein for the

purpose of cloning cDNA or genomic sequences which encode other parts or complete portions of the PAP genes described herein and therefore, these related sequences are encompassed within the present
5 invention.

In addition, isolated nucleic acid molecules of the invention include DNA molecules which comprise a sequence substantially different from those described above but which, due to the degeneracy of the genetic
10 code, still encode PAPs. Of course, the genetic code and species-specific codon preferences are well known in the art. Thus, it would be routine for one skilled in the art to generate the degenerate variants described above, for instance, to optimize codon
15 expression for a particular host (e.g., change codons in the human mRNA to those preferred by a bacterial host such as *E.coli* or plant host).

Nucleic acid molecules of the present invention may be in the form of RNA, such as mRNA, or in the
20 form of DNA, including, for instance, cDNA and genomic DNA obtained by cloning or produced synthetically. The DNA may be double-stranded or single-stranded. Single-stranded DNA or RNA may be the coding strand, also known as the sense strand, or it may be the non-
25 coding strand, also referred to as the antisense strand.

By "isolated" nucleic acid molecule(s) is intended a nucleic acid molecule, DNA or RNA, which has been removed from its native environment. For
30 example, recombinant DNA molecules contained in a vector are considered isolated for the purposes of the present invention. Further examples of isolated DNA molecules include recombinant DNA molecules maintained in heterologous host cells or purified (partially or
35 substantially) DNA molecules in solution. Isolated

RNA molecules include *in vivo* or *in vitro* RNA transcripts of the DNA molecules of the present invention. Isolated nucleic acid molecules according to the present invention further include such
5 molecules produced synthetically.

The present invention is further directed to nucleic acid molecules encoding portions or fragments of the nucleotide sequences described herein. Fragments include portions of the nucleotide sequences
10 of at least 10 contiguous nucleotides in length selected from any two integers, one of which representing a 5' nucleotide position and a second of which representing a 3' nucleotide position, where the first nucleotide for each nucleotide sequence is
15 position 1. That is, every combination of a 5' and 3' nucleotide position that a fragment at least 10 contiguous nucleotide bases in length or any integer between 10 and the length of an entire nucleotide sequence minus 1.

Further, the invention includes polynucleotides comprising fragments specified by size, in nucleotides, rather than by nucleotide positions. The invention includes any fragment size, in contiguous nucleotides, selected from integers between 1- and the
25 entire length of an entire nucleotide sequence minus 1. Preferred sizes include 20-50 nucleotides, 50-300 nucleotides useful as primers and probes. Regions from which typical sequences may be derived include but are not limited to, for example, regions encoding
30 specific epitopes or domains within said sequence, for example, the PBR binding domain extending in SEQ ID NO:1, 2, 3, 4, and 5, potential myristoylation sites at positions 262-267 and 271-276 of SEQ ID NO:7 and five PKC phosphorylation sites at 395-396, 113-115,
35 255-257, 280-282, 331-333, and 339-341 of SEQ ID NO:7,

an Acyl-Co-A site at position 24-108 of SEQ ID NO:7, a nuclear localization domain at position 150-167 of SEQ ID NO:7, a troponin site at position 98-247 of SEQ ID NO:7 and an HSP90 domain at position 126-155 of SEQ ID NO:7, among others.

In another aspect, the invention provides isolated nucleic acid molecules comprising polynucleotides which hybridize under stringent hybridization conditions to a polynucleotide sequence of the present invention described above, or a specified fragment thereof. By "stringent hybridization conditions" is intended overnight incubation at 42°C in a solution comprising: 50% formamide, 5X SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5X Denhardt's solution, 10% dextran sulfate, and 20 g/ml denatured sheared salmon sperm DNA, followed by washing the filters in 0.1X SSC at about 65°C.

The sequences encoding the polypeptides of the present invention or portions thereof may be fused to other sequences which provide additional functions known in the art such as a marker sequence, or a sequence encoding a peptide which facilitates purification of the fused polypeptide, peptides having antigenic determinants known to provide helper T-cell stimulation, peptides encoding sites for post-translational modifications, or amino acid sequences which target the fusion protein to a desired location, e.g. a heterologous leader sequence.

The present invention further relates to variants of the nucleic acid molecules of the present invention, which encode portions, analogs or derivatives of the PAPs of the present invention. A variant may occur naturally, such as a natural allelic variant. By an "allelic variant" is intended one of

several alternate forms of a gene occupying a given locus of a chromosome of an organism. Non-naturally occurring variants may be produced by known mutagenesis techniques. Such variants include those produced by
5 nucleotide substitution, deletion, or addition of one or more nucleotides in the coding or noncoding regions or both. Alterations in the coding regions may produce conservative or nonconservative amino acid substitutions, deletions, or additions. Especially
10 preferred among these are silent substitutions, additions, and deletions which do not alter the properties and activities of PAPs polypeptides disclosed herein or portions thereof. Also preferred in this regard are conservative substitutions.

15 Nucleic acid molecules with at least 90-99% identity to a nucleic acid identified above is another aspect of the present invention. These nucleic acids are included irrespective of whether they encode a polypeptide having PAP activity. By "a polypeptide
20 having PAP activity" is intended polypeptides exhibiting activity similar, but not identical, to an activity of the PAP of the invention, as measured in the assays described below. The biological activity or function of the polypeptides of the present
25 invention are expected to be similar or identical to polypeptides from other organisms that share a high degree of structural identity/similarity.

In another embodiment, the present invention relates to a recombinant DNA molecule that includes a
30 vector and a DNA sequence as described above. The vector can take the form of a plasmid, phage, cosmid, YAC, eukaryotic expression vector such as a DNA vector, *Pichia pastoris*, or a virus vector such as for example, baculovirus vectors, retroviral vectors or
35 adenoviral vectors, and others known in the art. The

cloned gene may optionally be placed under the control of (i.e., operably linked to) certain control sequences such as promoter sequences, or sequences which may be inducible and/or cell type-specific.

- 5 Suitable promoters will be known to a person with ordinary skill in the art. The expression construct will further contain sites for transcription initiation, termination and, in the transcribed region, a ribosome binding site for translation.
- 10 Among the vectors preferred for use include pGBT9, PGAD10 (Clonetech), PSVzeo (Invitrogen), pBlueScript (Stratagene), pCMV5 (Invitrogen), pCRII (Invitrogen) to name a few.

- Introduction of the construct into the host cell
- 15 can be effected by calcium phosphate transfection, electroporation, infection, and other methods known in the art and described in standard laboratory manuals such as Current Protocols in Molecular Biology, Ausubel, F. M. et al. (Eds), Wiley & Sons, Inc. All
- 20 documents cited herein supra and infra are hereby incorporated in their entirety by reference thereto.

- In a further embodiment, the present invention relates to host cells stably transformed or transfected with the above-described recombinant DNA
- 25 constructs. The host cell can be prokaryotic (for example, bacterial), lower eukaryotic (for example, yeast or insect) or higher eukaryotic (for example, all mammals, including but not limited to rat and human). Both prokaryotic and eukaryotic host cells
- 30 may be used for expression of desired coding sequences when appropriate control sequences which are compatible with the designated host are used. Among prokaryotic hosts, *E. coli* is most frequently used. Expression control sequences for prokaryotes include
- 35 promoters, optionally containing operator portions,

and ribosome binding sites. Transfer vectors compatible with prokaryotic hosts are commonly derived from, for example, pBR322, a plasmid containing operons conferring ampicillin and tetracycline resistance, and the various pUC vectors, which also contain sequences conferring antibiotic resistance markers. These markers may be used to obtain successful transformants by selection. Please see e.g., Maniatis, Fritsch and Sambrook, Molecular Cloning: A Laboratory Manual (1982) or DNA Cloning, Volumes I and II (D. N. Glover ed. 1985) for general cloning methods. The DNA sequence can be present in the vector operably linked to a sequence encoding an IgG molecule, an adjuvant, a carrier, or an agent for aid in purification of PAPs, such as glutathione S-transferase, or a series of histidine residues also known as a histidine tag. The recombinant molecule can be suitable for transfecting plant cells or eukaryotic cells, for example, mammalian cells and yeast cells in culture systems. *Saccharomyces cerevisiae*, *Saccharomyces carlsbergensis*, and *Pichia pastoris* are the most commonly used yeast hosts, and are convenient fungal hosts. Control sequences for yeast vectors are known in the art. Mammalian cell lines available as hosts for expression are known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC), such as HEK293 cells, and NIH 3T3 cells, MA10 Leydig cells, mouse C6 glioma cells, Y1 adrenal cells, and breast cancer cell lines such as MDA-231, MCF-7, to name a few. Suitable promoters are also known in the art and include viral promoters such as that from SV40, Rous sarcoma virus (RSV), adenovirus (ADV), bovine papilloma virus (BPV), and cytomegalovirus

(CMV). Mammalian cells may also require terminator sequences and poly A addition sequences; enhancer sequences which increase expression may also be included, and sequences which cause amplification of the gene may also be desirable. These sequences are known in the art. The transformed or transfected host cells can be used as a source of DNA sequences described above. When the recombinant molecule takes the form of an expression system, the transformed or transfected cells can be used as a source of the protein described below.

In another embodiment, the present invention relates to the PAP polypeptides described above or any allelic variation thereof which is immunologically identifiable with the polypeptides.

A polypeptide or amino acid sequence derived from the amino acid sequences mentioned above, refers to a polypeptide having an amino acid sequence identical to that of a polypeptide encoded in the sequence, or a portion thereof wherein the portion consists of at least 2-5 amino acids, and more preferably at least 8-10 amino acids, and even more preferably at least 11-15 amino acids, or which is immunologically identifiable with a polypeptide encoded in the sequence.

A recombinant or derived polypeptide is not necessarily translated from a designated nucleic acid sequence; it may be generated in any manner, including for example, chemical synthesis, or expression of a recombinant expression system. In addition the polypeptide can be fused to other proteins or polypeptides which increase its antigenicity, such as adjuvants for example.

As noted above, the methods of the present invention are suitable for production of any

polypeptide of any length, via insertion of the above-described nucleic acid molecules or vectors into a host cell and expression of the nucleotide sequence encoding the polypeptide of interest by the host cell.

5 Introduction of the nucleic acid molecules or vectors into a host cell to produce a transformed host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction,

10 infection or other methods. Such methods are described in many standard laboratory manuals, such as Davis et al., Basic Methods In Molecular Biology (1986). Once transformed host cells have been obtained, the cells may be cultivated under any

15 physiologically compatible conditions of pH and temperature, in any suitable nutrient medium containing assimilable sources of carbon, nitrogen and essential minerals that support host cell growth. Recombinant polypeptide-producing cultivation

20 conditions will vary according to the type of vector used to transform the host cells. For example, certain expression vectors comprise regulatory regions which require cell growth at certain temperatures, or addition of certain chemicals or inducing agents to

25 the cell growth medium, to initiate the gene expression resulting in the production of the recombinant polypeptide. Thus, the term "recombinant polypeptide-producing conditions," as used herein, is not meant to be limited to any one set of cultivation

30 conditions. Appropriate culture media and conditions for the above-described host cells and vectors are well-known in the art. Following its production in the host cells, the polypeptide of interest may be isolated by several techniques. To liberate the

35 polypeptide of interest from the host cells, the cells

are lysed or ruptured. This lysis may be accomplished by contacting the cells with a hypotonic solution, by treatment with a cell wall-disrupting enzyme such as lysozyme, by sonication, by treatment with high
5 pressure, or by a combination of the above methods. Other methods of bacterial cell disruption and lysis that are known to one of ordinary skill may also be used.

Following disruption, the polypeptide may be
10 separated from the cellular debris by any technique suitable for separation of particles in complex mixtures. The polypeptide may then be purified by well known isolation techniques. Suitable techniques for purification include, but are not limited to,
15 ammonium sulfate or ethanol precipitation, acid extraction, electrophoresis, immunoadsorption, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography,
20 immunoaffinity chromatography, size exclusion chromatography, liquid chromatography (LC), high performance LC (HPLC), fast performance LC (FPLC), hydroxylapatite chromatography and lectin chromatography.

25 The recombinant polypeptide or fusion protein can be used, detectably labeled and unlabeled, as a diagnostic tool for the detection of PAPs or for the detection and measurement of PBR. Additionally, these polypeptides can be used in a method for modulating
30 PBR expression. In addition, the recombinant protein can be used as a therapeutic agent to reduce cell death and/or increase cell proliferation via its effect on PBR function. The transformed host cells can be used to analyze the effectiveness of drugs and
35 agents which modulate PBR function, expression or

targeting via their effect on the expression or function of PAPs, such as host proteins or chemically derived agents or other proteins which may interact with the cell to alter the PAP function or expression, thereby modulating PBR function, expression or localization.

In another embodiment, the present invention relates to monoclonal or polyclonal antibodies specific for the above-described recombinant proteins (or polypeptides). For instance, an antibody can be raised against a peptide described above, or against a portion thereof of at least 10 amino acids, preferably, 11-15 amino acids. Persons with ordinary skill in the art using standard methodology can raise monoclonal and polyclonal antibodies to the protein (or polypeptide) of the present invention, or a unique portion thereof. Material and methods for producing antibodies are well known in the art (see for example Goding, in, Monoclonal Antibodies: Principles and Practice, Chapter 4, 1986).

The amount of PAP expression can be detected at several levels. Using standard methodology well known in the art, assays for the detection and quantitation of PAP RNA can be designed, and include northern hybridization assays, *in situ* hybridization assays, and PCR assays, among others. Please see e.g., Maniatis, Fitsch and Sambrook, Molecular Cloning: A Laboratory Manual (1982) or DNA Cloning, Volumes I and II (D. N. Glover ed. 1985), or Current Protocols in Molecular Biology, Ausubel, F. M. et al. (Eds), Wiley & Sons, Inc. for general description of methods for nucleic acid hybridization. Polynucleotide probes for the detection of PAP RNA can be designed from the sequences described in SEQ ID NO: 1-9. For example,

RNA isolated from samples can be coated onto a surface such as a nitrocellulose membrane and prepared for northern hybridization. In the case of *in situ* hybridization of biopsy samples for example, the
5 tissue sample can be prepared for hybridization by standard methods known in the art and hybridized with polynucleotide sequences which specifically recognize PAP RNA. The presence of a hybrid formed between the sample RNA and the polynucleotide can be detected by
10 any method known in the art such as radiochemistry, or immunochemistry, to name a few.

One of skill in the art may find it desirable to prepare probes that are fairly long and/or encompass regions of the amino acid sequence which would have a
15 high degree of redundancy in the corresponding nucleic acid sequences. In other cases, it may be desirable to use two sets of probes simultaneously, each to a different region of the gene. While the exact length of any probe employed is not critical, typical probe
20 sequences are no greater than 500 nucleotides, even more typically they are no greater than 250 nucleotides; they may be no greater than 100 nucleotides, and also may be no greater than 75 nucleotides in length. Longer probe sequences may be
25 necessary to encompass unique polynucleotide regions with differences sufficient to allow related target sequences to be distinguished. For this reason, probes are preferably from about 10 to about 100 nucleotides in length and more preferably from about
30 20 to about 50 nucleotides.

The DNA sequence of PAPs can be used to design primers for use in the detection of PAPs using the polymerase chain reaction (PCR) or reverse transcription PCR (RT-PCR). The primers can
35 specifically bind to the PAP cDNA produced by reverse

transcription of PAP RNA, for the purpose of detecting the presence, absence, or quantifying the amount of PAP by comparison to a standard. The primers can be any length ranging from 7-40 nucleotides, preferably 10-15 nucleotides, most preferably 18-25 nucleotides homologous or complementary to a region of the PAP sequence. Reagents and controls necessary for PCR or RT-PCR reactions are well known in the art. The amplified products can then be analyzed for the presence or absence of PAP sequences, for example by gel fractionation, by radiochemistry, and immunochemical techniques. This method is advantageous since it requires a small number of cells. Once PAP is detected, a determination whether the cell is overexpressing or underexpressing PAP can be made by comparison to the results obtained from a normal cell using the same method. For example, increased PAP7 RNA levels correlate with PBR expression levels, especially in steroidogenic cells, wherein, an increase in steroidogenic capability of the cells correlates with an increase in PBR and PAP7 RNA.

In another embodiment, the present invention relates to a diagnostic kit for the detection of PAP RNA in cells, said kit comprising a package unit having one or more containers of PAP oligonucleotide primers for detection of PAP by PCR or RT-PCR or PAP polynucleotides for the detection of PAP RNA in cells by *in situ* hybridization or northern analysis, and in some kits including containers of various reagents used for the method desired. The kit may also contain one or more of the following items: polymerization enzymes, buffers, instructions, controls, detection labels. Kits may include containers of reagents mixed together in suitable proportions for performing the

methods in accordance with the invention. Reagent containers preferably contain reagents in unit quantities that obviate measuring steps when performing the subject methods.

5 In a further embodiment, the present invention provides a method for identifying and quantifying the level of PAP present in a particular biological sample. Any of a variety of methods which are capable of identifying (or quantifying) the level of PAP in a
10 sample can be used for this purpose.

Diagnostic assays to detect PAPs may comprise a biopsy or *in situ* assay of cells from an organ or tissue sections, as well as an aspirate of cells from a tumor or normal tissue. In addition, assays may be
15 conducted upon cellular extracts from organs, tissues, cells, urine, or serum or blood or any other body fluid or extract.

When assaying a biopsy, the assay will comprise, contacting the sample to be assayed with a PAP ligand,
20 natural or synthetic, or an antibody, polyclonal or monoclonal, which recognizes PAP, or antiserum capable of detecting PAP, and detecting the complex formed between PAP present in the sample and the PAP ligand or antibody added.

25 PAP ligands or substrates include for example, PBR, in addition to natural and synthetic classes of ligands and their derivatives which can be derived from natural sources such as animal or plant extracts.

PAP ligands or anti-PAP antibodies, or fragments
30 of ligand and antibodies capable of detecting PAP may be labeled using any of a variety of labels and methods of labeling for use in diagnosis and prognosis of disease associated with increased cell proliferation, such as cancer, or reduced cell death.
35 Examples of types of labels which can be used in the

present invention include, but are not limited to, enzyme labels, radioisotopic labels, non-radioactive isotopic labels, and chemiluminescent labels.

Examples of suitable enzyme labels include malate
5 dehydrogenase, staphylococcal nuclease, delta-5-steroid isomerase, yeast-alcohol dehydrogenase, alpha-glycerol phosphate dehydrogenase, triose phosphate isomerase, peroxidase, alkaline phosphatase, asparaginase, glucose oxidase, beta-galactosidase,
10 ribonuclease, urease, catalase, glucose-6-phosphate dehydrogenase, glucoamylase, acetylcholine esterase, etc.

Examples of suitable radioisotopic labels include
15 ^3H , ^{111}In , ^{125}I , ^{32}P , ^{35}S , ^{14}C , ^{57}To , ^{58}Co , ^{59}Fe , ^{75}Se , ^{152}Eu , ^{90}Y , ^{67}Cu , ^{21}Ci , ^{211}At , ^{212}Pb , ^{47}Sc , ^{109}Pd , ^{11}C , ^{19}F , ^{123}I , etc.

Examples of suitable non-radioactive isotopic labels include ^{157}Gd , ^{55}Mn , ^{162}Dy , ^{52}Tr , ^{46}Fe , etc.

Examples of suitable fluorescent labels include a
20 ^{152}Eu label, a fluorescein label, an isothiocyanate label, a rhodamine label, a phycoerythrin label, a phycodyanin label, an allophycocyanin label, a fluorescamine label, etc.

Examples of chemiluminescent labels include a
25 luminal label, an isoluminal label, an aromatic acridinium ester label, an imidazole label, an acridinium salt label, an oxalate ester label, a luciferin label, a luciferase label, etc.

Those of ordinary skill in the art will know of
30 other suitable labels which may be employed in accordance with the present invention. The binding of these labels to ligands and to antibodies or fragments thereof can be accomplished using standard techniques commonly known to those of ordinary skill in the art. Typical techniques are described by Kennedy, J. H., et
35 *al.*, 1976 (*Clin. Chim. Acta* 70:1-31), and Schurs, A. H.

W. M., et al. 1977 (*Clin. Chim Acta* 81:1-40).

Coupling techniques mentioned in the latter are the glutaraldehyde method, the periodate method, the dimaleimide method, and others, all of which are incorporated by reference herein.

The detection of antibodies (or fragments of antibodies) of the present invention can be improved through the use of carriers. Well-known carriers include glass, polystyrene, polypropylene, polyethylene, dextran, nylon, amylases, natural and modified celluloses, polyacrylamides, agaroses, and magnetite. The nature of the carrier can be either soluble to some extent or insoluble for the purposes of the present invention. The support material may have virtually any possible structural configuration so long as the coupled molecule is capable of binding to PAP. Thus, the support configuration may be spherical, as in a bead, or cylindrical, as in the inside surface of a test tube, or the external surface of a rod. Alternatively, the surface may be flat such as a sheet, test strip, etc. Those skilled in the art will note many other suitable carriers for binding monoclonal antibody, or will be able to ascertain the same by use of routine experimentation.

The ligands or antibodies, or fragments of antibodies or ligands of PAPs discussed above may be used to quantitatively or qualitatively detect the presence of PAP. Such detection may be accomplished using any of a variety of immunoassays known to persons of ordinary skill in the art such as radioimmunoassays, immunometric assays, etc. Using standard methodology well known in the art, a diagnostic assay can be constructed by coating on a surface (i.e. a solid support) for example, a microtitration plate or a membrane (e.g.

nitrocellulose membrane), antibodies specific for PAP or a portion of PAP, and contacting it with a sample from a person suspected of having a PAP related disease. The presence of a resulting complex formed between PAP in the sample and antibodies specific therefor can be detected by any of the known detection methods common in the art such as fluorescent antibody spectroscopy or colorimetry. A good description of a radioimmune assay may be found in Laboratory Techniques and Biochemistry in Molecular Biology. by Work, T.S., et al. North Holland Publishing Company, N.Y. (1978), incorporated by reference herein. Sandwich assays are described by Wide at pages 199-206 of Radioimmune Assay Method, edited by Kirkham and Hunter, E. & S. Livingstone, Edinburgh, 1970.

The diagnostic methods of this invention can be predictive of diseases involving PBR including gallstones, atherosclerosis, Niemann-Pick C, Sitosterolemia, Dystrophy, Tumor proliferation (tumorigenesis), Schnyder's corneal crystalline dystrophy. Brain disorders include cholesterol metabolism and Alzheimer's disease, Tellurium toxicity, Smith-Lemli-Opitz syndrome, myelinization, developmental abnormalities and demyelization : Charcot-Marie-Tooth disease; Pelizaeus-Merzbacher disease, Multiple sclerosis, SLA, to name a few. Alternatively, the methods and compositions may be useful as prophylactic treatment, or in screening for compounds effective in prophylactic treatments.

The recombinant protein can be used to identify inhibitors or activators of a PAP activity which allows the identification of drugs or agents which modulate PBR activity. Using an assay as described below in the Examples, or for example, introducing a

drug or agent to a cell expressing a PAP and detecting
a increase or decrease in the level of PAP RNA or
protein, natural and synthetic agents and drugs can be
discovered which result in a reduction or elimination,
5 or increase in a PAP activity. Knowledge of the
mechanism of action of the inhibitor or activator is
not necessary as long as a decrease or increase in the
activity of a PAP is detected. Inhibitors may include
agents or drugs which either bind or sequester the
10 PAP's substrate(s), such as PBR, or cofactor(s), or
inhibit PAP itself, directly, for example by
irreversible binding of the agent or drug to the PAP,
or indirectly, for example by introducing an agent
which binds the competes with PAP binding to its
15 substrate. Activators may include cofactors necessary
for proper PAP function or agents which allow a higher
turnover rate of binding or release of the PAP to/from
PBR or the particular PAP substrate. Agents or drugs
related to this invention may result in partial or
20 complete inhibition or various degrees of activation
of PAP which may or may not result in modulation of
PBR function. Inhibitors or activators of PAP
activity may be used in the treatment or amelioration
of conditions such as stress, cancer,
25 neurodegenerative disorders, i.e. stroke, Alzheimer's,
developmental disorders, infertility, and immune
disorders.

Agents which decrease the level of PAP (i.e. in a
human or an animal) or reduce or inhibit PAP activity
30 may be used in the therapy of any disease associated
with the elevated levels of PAP. Similarly, agents
which increase the level of PAP or activate PAP
activity may be used in the therapy of any disease
associated with reduced levels of PAP. An increase or
35 decrease in the level of PAP is determined when the

change in the level of PAP is about 2-3 fold higher or lower than the level of PAP in the normal cell, up to about 10-100 fold higher or lower than the amount of PAP in a normal cell. Agents which decrease PAP RNA include, but are not limited to, one or more ribozymes capable of digesting PAP RNA, or antisense oligonucleotides capable of hybridizing to PAP RNA such that the translation of PAP is inhibited or reduced resulting in a decrease in the level of PAP. These agents can be administered as DNA, as DNA entrapped in proteoliposomes containing viral envelope receptor proteins (Kanoda, Y. et al., 1989, *Science* 243, 375) or as part of a vector which can be expressed in the target cell such that the DNA or RNA is made. Vectors which are expressed in particular cell types are known in the art, for example, for the mammary gland, please see Furth, (1997) (*J. Mammary Gland Biol. Neopl.* 2, 373) for examples of conditional control of gene expression in the mammary gland. Alternatively, the DNA can be injected along with a carrier. A carrier can be a protein such as a cytokine, for example interleukin 2, or polylysine-glycoprotein carrier. Such carrier proteins and vectors and methods of using same are known in the art. In addition, the DNA could be coated onto tiny gold beads and said beads introduced into the skin with, for example, a gene gun (Ulmer, J. B. et al., 1993, *Science* 259, 1745).

Alternatively, antibodies, or compounds capable of reducing or inhibiting PAP activity, that is reducing or inhibiting either the expression, production or activity of PAP, such as antagonists, can be provided as an isolated and substantially purified protein, or as part of an expression vector capable of being expressed in the target cell such

that the PAP-reducing or inhibiting agent is produced. Similarly, compounds capable of increasing or activating PAP activity, that is increasing or activating either the expression, production, or activity of PAP, such as agonists, can be provided as an isolated and substantially purified protein, or as part of an expression vector capable of being expressed in the target cell such that the PAP-elevating or activating agent is produced. In addition, factors which affect the stability of the protein, and co-factors such as various ions, i.e. Ca^{2+} (Calvo, D. J. and Medina, J. H., 1993, *J. Recept. Res.* 13:975-987), or anions, such as halides or anion channel blockers such as DIDS (4,4'-diisothiocyanostilbene-2,2'-disulfonic acid), an ion transport blocker (Skolnick, P., 1987, *Eur. J. Pharmacol.* 133:205-214), or factors which affect the stability of PBR such as lipids, for example, the phospholipids phosphatidylserine and phosphatidylinositol whereby the presence of the phospholipids is required for receptor activity (Moynagh, P. N. and Williams, D.C., 1992, *Biochem. Pharmacol.* 43:1939-1945) can be administered to modulate the expression and function of the PAPs and PBR. These formulations can be administered by standard routes. In general, the combinations may be administered by the topical, transdermal, intraperitoneal, oral, rectal, or parenteral (e.g. intravenous, subcutaneous, or intramuscular) route. In addition, PAP-inhibiting or PAP-activating compounds may be incorporated into biodegradable polymers being implanted in the vicinity of where drug delivery is desired, for example, at the site of a tumor or implanted so that the PAP-inhibiting or PAP-activating compound is slowly released systemically.

The biodegradable polymers and their use are described, for example, in detail in Brem *et al.* (1991) *J. Neurosurg.* 74, 441-446. These compounds are intended to be provided to recipient subjects in an amount sufficient to effect the inhibition of PAP. Similarly, agents which are capable of negatively or positively effecting the expression, production, stability or function of PAP, are intended to be provided to recipient subjects in an amount sufficient to result in the desired effect. An amount is said to be sufficient to "effect" the inhibition or induction of PAP if the dosage, route of administration, etc. of the agent are sufficient to influence such a response.

The PAPs identified in this application were discovered due to their ability to associate with PBR, and may play a role in the proper targeting, function, expression, or stability of PBR. Therefore, a method for inhibiting or reducing PBR function, or altering the localization of PBR, would include a method for dissociating PAPs from the receptor. This is possible using agents which block the site on PBR at which these PAPs associate with PBR, or alternatively, blocking the site on the PAPs which is involved in PBR-association. Such agents would include antibodies or antagonists which recognize such sites or which alter the conformation of these sites such that PAP and PBR association is inhibited or eliminated. Agents which decrease the level of PBR (i.e. in a human or an animal) or reduce or inhibit PBR activity may be used in the therapy of any disease associated with the elevated levels of PBR such as metastatic cancer, for example breast cancer, or diseases associated with increased cell proliferation or increased cholesterol transport into the cell. An increase in the level of PBR is determined when the

level of PBR in a tumor cell is about 2-3 times the level of PBR in the normal cell, up to about 10-100 times the amount of PBR in a normal cell.

Antibodies or compounds capable of reducing or inhibiting the association between PBR and PAPs can be provided as an isolated and substantially purified protein, or as part of an expression vector capable of being expressed in the target cell such that the PBR-PAP-association reducing or inhibiting agent is produced. These formulations can be administered by standard routes. In general, the combinations may be administered by the topical, transdermal, intraperitoneal, oral, rectal, or parenteral (e.g. intravenous, subcutaneous, or intramuscular) route. In addition, these compounds may be incorporated into biodegradable polymers being implanted in the vicinity of where drug delivery is desired, for example, at the site of a tumor or implanted so that the compound is slowly released systemically. The biodegradable polymers and their use are described, for example, in detail in Brem *et al.* (1991) *J. Neurosurg.* 74, 441-446.

These compounds are intended to be provided to recipient subjects in an amount sufficient to effect the inhibition of PBR/PAP association.

In line with the function of PBR in cell proliferation, agents which stimulate the function of PBR by increasing the association of PAPs to PBR, may be used in the therapy of any disease associated with a decrease of PBR, or a decrease in cell proliferation, wherein PBR is capable of increasing such proliferation, e.g. developmental retardation. PBR has also been shown to be involved in cholesterol transport, therefore, an agent or drug which results in an increase in function or stability of PBR and its associated PAPs can be used to increase cholesterol

transport into cells. Diseases where cholesterol transport is deficient include lipoidal adrenal hyperplasia, and diseases where there is a requirement for increased production of compounds requiring cholesterol such as myelin and myelination including Alzheimer's disease, spinal chord injury, and brain development neuropathy [Snipes, G. and Suter, U. (1997) Cholesterol and Myelin. In: Subcellular Biochemistry, Robert Bittman (ed.), vol. 28, pp.173-204, Plenum Press, New York], to name a few.

In providing a patient with any agent which modulates the expression, function, targeting, or association of PAP or PBR as discussed above, the dosage of administered agent will vary depending upon such factors as the patient's age, weight, height, sex, general medical condition, previous medical history, etc. In general, it is desirable to provide the recipient with a dosage of agent which is in the range of from about 1 pg/kg to 10 mg/kg (body weight of patient), although a lower or higher dosage may be administered.

A composition is said to be "pharmacologically acceptable" if its administration can be tolerated by a recipient patient. Such an agent is said to be administered in a "therapeutically effective amount" if the amount administered is physiologically significant. An agent is physiologically significant if its presence results in a detectable change in the physiology of a recipient patient.

The compounds of the present invention can be formulated according to known methods to prepare pharmaceutically useful compositions, whereby these materials, or their functional derivatives, are combined in admixture with a pharmaceutically

acceptable carrier vehicle. Suitable vehicles and their formulation, inclusive of other human proteins, e.g., human serum albumin, are described, for example, in Remington's Pharmaceutical Sciences [16th ed., Osol, A. ed., Mack Easton PA. (1980)]. In order to form a pharmaceutically acceptable composition suitable for effective administration, such compositions will contain an effective amount of the above-described compounds together with a suitable amount of carrier vehicle.

Additional pharmaceutical methods may be employed to control the duration of action. Control release preparations may be achieved through the use of polymers to complex or absorb the compounds. The controlled delivery may be exercised by selecting appropriate macromolecules (for example polyesters, polyamino acids, polyvinyl, pyrrolidone, ethylenevinylacetate, methylcellulose, carboxymethylcellulose, or protamine sulfate) and the concentration of macromolecules as well as the method of incorporation in order to control release. Another possible method to control the duration of action by controlled release preparations is to incorporate the compounds of the present invention into particles of a polymeric material such as polyesters, polyamino acids, hydrogels, poly(lactic acid) or ethylene vinylacetate copolymers. Alternatively, instead of incorporating these agents into polymeric particles, it is possible to entrap these materials in microcapsules prepared, for example, interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly(methylmethacrylate)microcapsules, respectively, or in colloidal drug delivery systems, for example, liposomes, albumin microspheres, microemulsions,

nanoparticles, and nanocapsules or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences (1980).

5 The present invention also provides kits for use in the diagnostic or therapeutic methods described above. Kits according to this aspect of the invention may comprise one or more containers, such as vials, tubes, ampules, bottles and the like, which may
10 comprise one or more of the compositions of the invention.

 The kits of the invention may comprise one or more of the following components, one or more compounds or compositions of the invention, and one or more excipient, diluent, or adjuvant.

15 It will be readily apparent to one of ordinary skill in the relevant arts that other suitable modifications and adaptations to the methods and applications described herein are obvious and may be made without departing from the scope of the invention
20 or any embodiment thereof. Having now described the present invention in detail, the same will be more clearly understood by reference to the following examples, which are included herewith for purposes of illustration only and are not intended to be limiting
25 of the invention.

 The following Materials and Methods were used in the Examples described below.

Material and Methods

Materials

30 [α -³²P]dCTP (specific activity, 3000 Ci/mmol), [1,2,6,7-N-³H]progesterone (specific activity, 94.1 Ci/mmol) and ³H-1-(2-chlorophenyl)-N-methyl-N-(1-methyl-propyl)-3-isoquinolinecarboxamide (PK 11195) (specific activity, 86.9 Ci/mmol) were obtained from

NEN Life Science Products (Boston, MA). PK11195 was obtained from Research Biochemicals, Inc. (Natick, MA). Nitrocellulose (0.45 μ m) was from Hoeffer Scientific (San Francisco, CA). 22R

5 Hydroxycholesterol was purchased from Sigma. Restriction enzymes were from Stratagene (La Jolla, CA) and New England Biolabs (Beverly, MA). Cell culture supplies were purchased from Life Technologies, Inc. (Grand Island, NY). Tissue culture
10 plasticware was from Corning (Corning, NY). Electrophoresis reagents and materials were supplied from BioRad. All other chemicals used were of analytical grade and were obtained from various commercial sources.

15 Strains and media

The genotype of the *Saccharomyces cerevisiae* reporter strain HF7c is *MATa*, *ura3-52*, *his3-200*, *lys2-801*, *ade2-101*, *trp1-901*, *leu2-3, 112*, *gal4-542*, *gal80-538*, *LYS2::GAL-HIS3*, *URA3::(GAL4 17-mers)₃-CYC1-lacZ*
20 (CLONTECH, Palo Alto, CA). Yeast strains were grown at 30°C in standard liquid YPD medium or minimal SD synthetic medium with appropriate supplement amino acids (CLONTECH, Palo Alto, CA).

Plasmids and construction

25 The mouse PBR cDNA coding sequence was subcloned into pGBT9 (CLONTECH, Palo Alto, CA) at *EcoR I* and *BamH I* sites (pGBT-PBR). The fusion site was verified by sequencing. Functional fusion PBR protein, expressed in yeast cells, was verified by PBR ligand
30 binding assay. Mouse testis cDNA library was constructed in pGAD10 [LEU2, GAL4 (768-881)] (CLONTECH, Palo Alto, CA). Amplification of premade libraries was performed by growing the transformants on LB-agar-ampicillin and purifying the plasmids DNA

with QIAGEN Plasmid Giga kit (QIAGEN, Valencia, CA).
In the transfection experiments, PAP7 partial sequence
(including 192 Amino acids C-terminal sequence) was
inserted into pSVzeo vector (Invitrogen, Carlsbad ,
5 CA) at *EcoRI* and *BamHI* sites.

Yeast two-hybrid screening

The Clontech MATCHMAKER two-hybrid system was
applied in this study (detailed in manufacturer's
instruction book). Briefly, the yeast reporter host
10 strain HF7c was simultaneously cotransformed with both
pGBT-PBR and the mouse testis cDNA library in pGAD10
plasmid by using lithium acetate high-efficiency
method (Gietz, D. et al., 1992, *Nucleic Acids Res.* 20,
1425). HIS positive clones were further selected by
15 colony lift filter assay for β -galactosidase activity.
Plasmid DNA was rescued in *Escherichia coli* DH5 α from
yeast cells. Plasmids were retransformed into yeast
HF7c cells with plasmid pGBT-PBR to test for histidine
prototrophy and β -galactosidase activity (Clontech
20 manual). The cDNA inserts from the positive clones
were sequenced. The full length PAP7 cDNA was
obtained by using 5' and 3' RACE kit from kit
(CLONTECH, Palo Alto, CA).

Sequence analysis

25 The ABI PRISMTM dyes terminator cycle sequencing
ready reaction kit (PE Biosystems, Foster City , CA)
and an Applied Biosystems sequencer were used for
sequencing (Applied Biosystems, Foster City , CA) at
the Lombardi Cancer Center Sequencing Core Facility
30 (Georgetown University). DNA sequences were analyzed
by using Entrez and BLAST program against GeneBankTM
Database.

Cell culture transient transfection

MA-10 cells were grown in modified Waymouth's MB752/1 medium containing 15% horse serum, as described previously (Papadopoulos, V. et al., 1990, *J. Biol. Chem.* 265, 3772-3779). Mouse C6 glioma and mouse Y1 adrenal cortical cells were cultured in DMEM and DMEM F12 respectively, with 10% fetal bovine serum. MA10 cells were transiently transfected by electroporation (El Hefnawy, T. et al., 1996, *Mol. Cell Endocrinol.* 119, 207-217). Each Genepulser cuvette (0.4 cm-gap, BioRad, Hercules, CA) contained 8×10^6 cells in 350 μ l antibiotic-free complete Waymouth's growth medium (see above), plus 30 μ g plasmids DNA in 50 μ l of 0.1 x TE. Cells in electroporation cuvettes were electro-shocked at 330 V and at a capacitance of 950 μ Fd generated from Genepulser (BioRad, Hercules, CA). The cells were kept immediately on ice for 10 min before plated into 96 well plates.

Radioligand binding assays

^3H -1-(2-chlorophenyl)-N-methyl-N-(1-methyl-propyl)-3-isoquinolinecarboxamide (PK 11195) (NEN, Boston, MA) binding studies were performed as we previously described (Papadopoulos, V. et al., 1990, *supra*; Garnier, M. et al., 1994, *Molecular Pharmacology* 45, 201-211). The dissociation constant (Kd) and the number of binding sites (Bmax) were determined by Scatchard plot analysis of the data using the LIGAND program (Munson, P. J. and Rodbard, D., 1980, *Anal. Biochem.* 107, 220-239).

RNA (Northern) blot analysis

Total tissue and cellular RNA was isolated by the acid guanidinium thiocyanate-phenol-chloroform extraction method using RNA STAT60 reagent (Tel-Test

Inc., Friendswood, TX). RNA was separated by denature electrophoresis and transferred to Nytran membrane (Schleicher & Schuell Inc., Keene, NH). The RNA blots were hybridized with ^{32}P labeled PAP7 cDNA probe generated from random priming (Boehringer Mannheim, Indianapolis, IN). Autoradiography was performed by exposing Kodak X-Omat AR films (Eastman Kodak, Rochester, NY) to the blots at -80°C overnight.

Steroid biosynthesis

MA-10 cells were plated into 96-well plate at the density of 2.5×10^4 /well for overnight. The cells were stimulated with 50 ng/ml hCG in 0.2 ml/well serum-free medium for 2 hours. The culture medium was collected and tested for progesterone production by RIA. The assay was carried out by using anti-progesterone antisera (ICN, Costa Mesa, CA), following the conditions recommended by the manufacturer. The progesterone production was normalized by the amount of protein in each well. Radioimmunoassay data was analyzed using the software provided by Wallac (EG&G Wallac, Gaithersburg, MD).

Antibody generation and Western analysis

Rabbit anti-PAP7 antibody was prepared by sequential immunization with a peptide SSDEEEEEENVTCEEKAKKNANKP (SEQ ID NO:11) of PAP7 protein, which was coupled to KLH. PAP7 antibodies were purified by an affinity resin containing the same peptide immobilized onto agarose (Bethyl Laboratories, Montgomery, TX). MA10 cells were solubilized in sample buffer (25 mM Tris-HCl (pH6.8), 1% SDS, 5% β -mercaptoethanol, 1 mM EDTA, 4% glycerol, and 0.01% bromophenol blue), boiled for 5 min, and loaded onto a 15% SDS-PAGE minigel (MiniProtein II System, BioRad, Richmond, CA). Electrophoresis was performed at 25

mA/gel using a standard SDS-PAGE running buffer (25 mM Tris, 192 mM glycine, and 0.1% SDS). The proteins were electrophoretically transferred to nitrocellulose membrane (Schleicher & Schuell Inc., Keene, NH). The
5 membrane was incubated in blocking buffer (TTBS buffer (20 mM Tris/HCl, pH 7.5, 0.5M NaCl, and 0.05% Tween-20) containing 10% Carnation nonfat milk) at room temperature for 1 hour, followed by incubation with a primary antibody against PAP7 (1:2000) for 2 hours.
10 The membrane was washed with TTBS three times for 10 min each time. After one-hour incubation with the secondary antibody, goat anti-rabbit IgG conjugated with HRP (Signal Transduction), the membrane was washed with TTBS three times for 10 min each time.
15 Specific protein bands were detected by chemiluminescence using the Renaissance Kit (DuPont-New England Nuclear, Wilmington, DE) according to manufacturer's directions.

Immunocytochemistry

20 MA-10 cells were cultured on four-chambered SuperCell Culture Slides (Fisher Scientific, Pittsburgh, PA) and fixed with methanol at 4°C for 15 min. The fixed cells were incubated with PAP7 antibody (1:250 dilution) with or without PAP7 peptide
25 for 1 hour. After washing, the cells were incubated with HRP conjugated goat anti-rabbit secondary antibody (Transduction Lab, Lexington, KY) for 1 hour. PAP7 staining was visualized with peroxidase using AEC (3-amino-9-ethyl carbazole) as a chromogen to yield a
30 red reaction product. After counterstaining with hematoxylin, slides were dehydrated and permanently mounted.

Immunohistochemistry

Mouse tissues were freshly snapped in liquid nitrogen. Specimens were fixed in cold methanol right after sectioning for 5 min. The slides were then placed in a chamber containing 0.3% H₂O₂ solution in methanol for 20 min at room temperature to inhibit the endogenous peroxidase activity and then incubated in blocking solution (10% goat serum) (Zymed, South San Francisco, CA) for 15 min. Subsequently, the slides were incubated with anti-PAP7 antibody (1:250) for 2 h at room temperature, washed with water and PBS, incubated with HRP conjugated goat anti-rabbit secondary antibody for 1 hour at room temperature, and then washed with PBS. After treatment with AEC reagent for 1 hr at 37°C for color staining, the sections were counterstained with hematoxylin, dehydrated and permanently mounted.

Protein quantification and Statistical analysis

Proteins were quantified by dye-binding assay of Bradford (Bradford, M. M., 1976, *Anal. Biochem.* 72, 248-254) with bovine serum albumin as the standard. Statistical analysis was performed by ANOVA followed by the Student-Newman-Keuls test or the Dunnett multiple comparisons test using the Instat (v.2.04) package from GraphPad, Inc. (San Diego, CA).

Example 1

Isolation of PBR Associated Proteins

We have used the MATCHMAKER Two-Hybrid System from CLONTECH in order to clone genes whose products interact with PBR protein. GAL4 (1-147)-PBR fusion (plasmid pGBT9 + PBR) was used as a bait to screen a mouse MATCHMAKER testis cDNA library constructed into the pGAD10 two-hybrid vector. About 3x10⁶ transformants were tested, and five positive clones were obtained for their ability to interact with PBR.

Library plasmids from these transformants were rescued in *E. coli* strain DH5 α . Both the His⁺ phenotype and the expression of β -galactosidase were confirmed by a second-round transformation of strain HF7c carrying pGBT9-PBR (Table 1).

Table 1: Summary of yeast two hybrid screen of mouse testis library by PBR

Clone	His3	β -Galactosidase
activity		
PAP3	+	++
PAP7	+	++
PAP20	+	++
Positive control	+	+++

Plasmids from these positive clones were first analyzed by restriction enzyme digestion and followed by sequence analysis. Two clones were shown to be coded by single gene, which was an unknown gene and was named as PBR associated protein 7 (PAP7). The other three clones encoded different products. After 5' RACE and 3' RACE, PAP7 cDNA clone was completely sequenced on both strands (SEQ ID NO:2) and it encoded a 463-amino acid protein with a calculated molecular weight of about 52 kDa. A homology search in the Genbank database using the BLAST program showed that this is a new sequence, previously unidentified.

Example 2

PAP7 protein expression in MA-10 Leydig tumor cells

The total MA-10 cell protein extracts were analyzed by western blot using PAP7 antibody. This antibody specifically recognizes a 50 kDa-protein band (Fig. 1A). The PAP7 protein expression in MA-10 cell

was also checked by immunocytochemistry. PAP7 antibody specifically stained MA-10 cell, with the signal mostly localized in the cytoplasm (Figure 1B). This signal can be neutralized by PAP7 peptide, which was used to generate and purify this antibody.

Example 3

PAP7 cells and tissue expression by Dot and Northern Blot

By dot blot analysis, PAP7 was observed to be highly expressed in brain, eye, submax gland, testis, and ovary. Interestingly, PAP7 expression was at its highest level at early embryonic stage, and decreased before birth (Fig. 2A and 2B). Consistently, PAP7 mRNA was expressed in adrenal, brain, heart, liver, testis and ovarian tissues by Northern blot analysis. PAP7 had a 1 kb transcript which was only expressed in testis and a 3 Kb major mRNA transcript in the other tissues (Fig. 3A and 3B). PAP7 was also highly expressed in three cell lines, C6 glioma, MA-10 Leydig cells and Y1 adrenal cells, which have been widely used for studying steroid biosynthesis. All three cell lines expressed PAP7 transcript of the same molecular weight size as in normal tissues. The PAP7 expression level in these cell lines was proportionally correlated with their steroidogenic capability (Fig. 3A and 3B). The PBR mRNA expression level was also checked in these same tissues and cell lines. The PBR expression level was parallel with PAP7 mRNA expression pattern, especially in those three cell lines (Fig. 4A and 4B).

Example 4

PAP7 Cellular distribution

PAP7 protein expression in different tissues was checked by immunohistochemistry (Fig 5). PAP7 was

present in both Leydig and germ cells in testis (Figure 5C and 5D), in hippocampus and neuronal cells in brain (Figure 5E and 5F), in fasciculata cells in adrenal gland (Figure 5G and 5H), and in granulosa cells in ovary (Figure 5A and 5B). Liver and kidney expressed low level of PAP7 protein (data not shown). Each specimen was stained with PAP7 peptide neutralized antibody as a negative control. Subsequent in situ hybridization studies showed that PAP7 mRNA followed PAP-7 protein expression.

Example 5

The effect of PAP7 on steroid biosynthesis in MA-10 cell

PAP7 partial sequence including PBR binding domain was subcloned into pSVzeo mammalian expression vector. This pSVzeoPAP7 vector was transiently transfected into MA-10 cells. pSVzeo empty vector was also transfected into cells as control. The capability of steroid biosynthesis of both empty vector pSVzeo transfectants and pSVzeoPAP7 transfectants was checked by monitoring the progesterone production in response to hormonal (hCG) stimulation. PAP7 transfectants had significantly reduced the level of progesterone production in MA-10 cells as compared with pSVzeo vector transfectant at a dose and time dependent manner (Fig 6).

Discussion

In order to better understand the mechanism of how PBR regulates cholesterol transport activity in steroid biosynthesis, we performed yeast two-hybrid assay to identify the PBR associated protein(s). Mouse PBR cDNA was inserted into the pGBT9 vector to generate a GAL4 DNA-binding domain and PBR fusion

protein as bait. The receptor ligand binding activity of the fusion protein with PBR ligand, PK11195, was tested. Our result indicated that the fusion PBR protein expressed in yeast possessed the similar binding affinity as the native PBR protein (data not shown). PBR has been identified in various peripheral tissues (Gavish, M. and Weizman, R., 1997, *Clin. Neuropharmacol.* 20, 473-481) including testis. Testis is one of the important and very well studied tissues for steroidogenesis (Huhtaniemi, I. and Toppari, J., 1995, *Adv. Exp. Med. Biol.* 377, 33-54). The steroid biosynthesis in the mouse testicular Leydig cell and the role of PBR in this process are also well documented (Papadopoulos, V. et al., 1997, *Steroids* 62, 21-28; Papadopoulos, V. et al., 1998, *Endocr. Res.* 24, 479-487). Therefore, we applied mouse testis cDNA library in this two-hybrid screen study. The pGAD10 vector was used to generate a fusion protein of the GAL4 activation domain with a collection of random proteins in the fusion Balb/c mouse testis library. Through the yeast two hybrid screen, PAP7 was identified as one of the positive clones, which demonstrated its ability to interact with PBR (Table 1). Thus, we cloned the PAP7 cDNA coding for a mouse protein that interacted with PBR. Based on the database search, PAP7 is a novel gene product. Recently, PRAX-1 was reported as a new protein that specifically interacts with PBR (Galiege, S. et al., 1999, *J. Biol. Chem.* 274, 2938-2952). The only similarity is that both proteins contain glutamic-acid stretches. Part of PAP7 shares quite high homology with a *C. elegans* gene that has an unknown function (Wilson, R. et al., 1994, *Nature* 368, 32-38). In fact, cholesterol is required for *C. elegans* cell culture (Brenner, S., 1974, *Genetics* 77, 71-94).

Considering that PBR is involved in cholesterol transport and PBR gene is highly conservative in all type of organisms, this data suggests that PAP7 expression may be needed to meet basic requirements for cell survival and growth. PAP7 also shares some homologies with RALBP, a hydrophobic ligand-binding protein that functions in intracellular retinoid transport (Ozaki, K. et al., 1994, *J. Biol. Chem.* 269, 3838-3845).

By sequence motif analysis using Swiss-Port Prosite profile scan, PAP7 has fatty acylation (myristoylation) sites, Acyl-CoA-binding protein signature and PKC phosphorylation sites. Protein myristoylation enables protein to attach to the cellular membrane and thus take part in cell signaling (Casey, P.J., 1995, *Science* 268, 221-225; Boutin, J. A., 1997, *Cell Signal* 9, 15-35). PBR is a hydrophobic protein and tightly associated with the outer mitochondrial membrane. This property could enable PAP7 passing hormone stimulation signal to and interacting with PBR thus regulating PBR activity in cholesterol transport. Interestingly, Acyl-CoA-binding protein is the other name of PBR endogenous ligand, diazepam binding inhibitor (DBI) (Rose, T. M. et al., 1992, *Proc. Natl. Acad. Sci. U.S.A.* 89, 11287-11291; Costa, E. and Guidotti, A., 1991, *Life Sci.* 49, 325-344; Suk, K. et al., 1999, *Biochim. Biophys. Acta* 1454, 126-131). This information suggests that PAP7 may coordinate with other PBR endogenous ligands to fulfill its function. The fact that PAP7 has the potential protein kinase phosphorylation sites raises another possibility that PBR could be regulated by hormone stimulation through the interaction with PAP7 protein.

The distribution and expression of PAP7 were examined in major mouse tissues such as brain, testis, ovary, adrenal, and kidney, as well as some cell lines. The PAP7 expressing pattern is similar to the broader expression profile of PBR. According to the previous studies (Papadopoulos, V. et al., 1998, supra), glucocorticoids are produced by zona fasciculata cells in adrenal. In the ovary, corpus luteum, where the granulosa cells are located, secretes progesterone. In addition, the testicular Leydig cells are able to produce testosterone. Since PAP7 is highly expressed in major steroidogenic tissues and is more concentrated in these steroid producing cells, PAP7 may be involved in steroid biosynthesis or the regulation of steroid formation by changing the formation or the conformation of the PBR complex. Mouse C6 glioma cells, MA-10 Leydig cells and Y1 adrenal cortical cells are popular cell models selected for studying steroid biosynthesis. PAP7 expression is proportionally correlated to the PBR expression in these cell lines. Additionally, both PBR and PAP7 expression level in these cell lines are parallel with their steroidogenic capability, which also suggests that PAP7 may be involved in steroid biosynthesis through a PBR pathway. A small PAP7 transcript was expressed only in testis, a phenomenon observed for other genes expressed in testis (Zhang, F. P. et al., 1997, *Endocrinology* 138, 2481-2490; Mauduit, C. et al., 1999, *J. Biol. Chem.* 274, 770-775). The immunostaining in the testis other than in Leydig cells could represent the expression of the smaller transcript.

The PAP7 protein is expressed in MA-10 cells and most of the staining is localized in the cytoplasm. The study of PAP7 subcellular distribution is ongoing,

the results of which may provide more detailed information about the interaction between PBR and PAP7. Since PBR knockout mice die in uterus indicates an essential role for PBR essential in mouse embryonic development. Interestingly, PAP7 mRNA is highly expressed during mouse early embryonic development. This result may suggest that PAP7, associated with PBR, could play an important role during early mouse development. This further implies that PBR may have new functions beyond steroidogenesis. Overexpression of PAP7 fragment including its PBR binding domain significantly inhibited the progesterone formation stimulated by saturating concentrations of hCG (50 ng/ml) in MA-10 cells. According to previous studies, progesterone production of these cells represented the index of steroid biosynthesis (Freeman, D. A., 1987, *Endocrinology* 120, 124-132; Garnier, M. et al., 1994, *J. Biol. Chem.* 269, 22105-22112). Based on the inhibitory manner, we assume that the overexpressed PAP7 fragment might act as a competitor of the native PAP7 in MA-10 cells and competitively bind to PBR. We believe that the transfected PAP7 fragment having only the PBR binding domain and is not fully functional as a native PAP7, however, it competitively prevents PBR from interacting with the endogenous PAP7 and thereby blocks the normal function of PBR.

In conclusion, the results presented herein suggest that the identified PAP7 is involved in the regulation of the PBR function, serving as an endogenous ligand or allosteric modulator of the receptor.

Considering the findings that i) PBR is a channel/transporter of cholesterol, ii) PBR is the target of environmental antisteroidogenic hazards, and (iii) PBR is involved in breast cancer aggression and

tumor cell proliferation (Hardwick, M. et al., 1999, *Cancer Res.* 59, 831-842), we believe that the identification and characterization of PAP7 will greatly contribute to the understanding of the role of PBR in steroidogenesis and even in more general areas.

10

15

20

25

30

35

What is claimed is:

1. An isolated PBR-associated protein (PAP) DNA fragment or any portion thereof.
2. An isolated and purified DNA fragment which
5 encodes a PBR-associated protein.
3. An isolated and purified DNA fragment which encodes a peptide of PBR-associated protein, said DNA fragment comprising a sequence specified in any of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, and
10 SEQ ID NO:5, or polynucleotide fragment of said sequence comprising at least 30 nucleotides.
4. An isolated and purified DNA fragment which encodes a peptide of PBR-associated protein, said DNA fragment comprising a sequence specified in Genbank
15 Accession no. AF022770, or GenBank Accession no. AF020338, or a polynucleotide fragment of said sequence comprising at least 30 nucleotides.
5. An isolated and purified PAP7 DNA fragment according to claim 2 which encodes 463 amino acids of
20 PAP7 or a natural variant or synthetic variant thereof encoding PAP7, or a peptide fragment thereof comprising at least 10 amino acids.
6. An isolated and purified PAP8 DNA fragment according to claim 2 which encodes 190 amino acids of
25 PAP8 or a natural variant or synthetic variant thereof encoding PAP8, or a peptide fragment thereof comprising at least 10 amino acids.
7. An isolated and purified PAP15 DNA fragment according to claim 2 which encodes 164 amino acids of
30 PAP15 or a natural variant or synthetic variant thereof encoding PAP15, or a peptide fragment thereof comprising at least 10 amino acids.
8. An isolated and purified PAP20 DNA fragment according to claim 2 which encodes 196 amino acids of
35 PAP20 or a natural variant or synthetic variant

thereof encoding PAP20, or a peptide fragment thereof comprising at least 10 amino acids.

9. A recombinant DNA construct comprising:

(i) a vector, and

5 (ii) the PAP DNA fragment of claim 1.

10. A recombinant DNA construct comprising:

(i) a vector, and

(ii) the PAP DNA fragment of claim 3.

11. A recombinant DNA construct according to
10 claim 10, wherein said vector is an expression vector.

12. The recombinant DNA construct according to
claim 10, wherein said vector is a prokaryotic vector.

13. The recombinant DNA construct according to
claim 10, wherein said vector is a eukaryotic vector.

15 14. A host cell transformed with a recombinant
DNA construct according to claim 10.

15. A host cell according to claim 14, wherein
said cell is prokaryotic.

20 16. A host cell according to claim 14, wherein
said cell is eukaryotic.

17. A method for producing PAP peptide which
comprises culturing the cells according to either
claim 15 or 16, under conditions such that said DNA
fragment is expressed and said PAP peptide is thereby
25 produced.

18. An isolated recombinant PAP produced by the
method of claim 17.

19. A PAP7 polypeptide comprising the amino acid
sequence specified in SEQ ID NO:7 or a portion thereof
30 of at least 5 amino acids.

20. A PAP8 polypeptide comprising the amino acid
sequence specified in SEQ ID NO:8 or a portion thereof
of at least 5 amino acids.

21. A PAP15 polypeptide comprising the amino acid sequence specified in SEQ ID NO:9 or a portion thereof of at least 5 amino acids.

22. A PAP20 polypeptide comprising the amino acid
5 sequence specified in SEQ ID NO:10 or a portion thereof of at least 5 amino acids.

23. A method for detecting a PAP in a sample chosen from the group consisting of: PAP7 , PAP8, PAP15, PAP20, said method comprising

- 10 (i) contacting said sample with antibodies which recognize said PAP; and
(ii) detecting the presence or absence of a complex formed between PAP and antibodies specific therefor.

15 24. A method for detecting a PBR-associated protein, said method comprising the two hybrid assay.

25. An antibody to a peptide having the amino acid sequence specified in SEQ ID NO:6, 7, 8 and 9, or any portion thereof.

20 26. A PAP7 antibody to a peptide comprising the amino acid sequence specified in SEQ ID NO:11.

27. A method for detecting agents or drugs which reduce or eliminate PAP activity, said method comprising:

- 25 (i) delivering a recombinant DNA construct according to claim 10 into a cell such that PAP is produced in said cell;
(ii) adding at least one drug or agent to said cell alone or in combination; and,
30 (iii) detecting PAP activity in said cell in the presence of said agent or drug and comparing it to a control which did not receive said drug or agent wherein a decrease in PAP activity as compared to control indicates an drug or agent which reduces or
35 eliminates PAP activity.

28. A method for detecting agents or drugs which promote PAP activity, said method comprising:

(i) delivering a recombinant DNA construct according to claim 10 into a cell such that PAP is produced in said cell;

(ii) adding at least one drug or agent to said cell alone or in combination; and,

(iii) detecting whether or not said drug or agent stimulates PAP activity by measuring PAP activity in said cell and comparing it to a control which did not receive said drug or agent wherein an increase in the activity of said PAP in said cell as compared to control indicates a stimulatory drug or agent.

29. An agent or drug capable of inhibiting PAP activity.

30. An agent or drug capable of promoting PAP activity.

31. A therapeutic compound comprising said agent or drug according to claim 29 for use in a disease wherein a decrease or elimination of PAP activity is beneficial.

32. A therapeutic compound comprising said agent or drug according to claim 30 for use in a disease wherein an increase o PAP activity is beneficial.

33. A method for detecting at least one PAP selected from the group consisting of PAP7, PAP8, PAP15, and PAP20 in a sample using the polymerase chain reaction.

34. A diagnostic kit for detecting RNA/cDNA of at least one PAP chosen from the group consisting of PAP7, PAP8, PAP15 and PAP20, in a sample comprising primers or oligonucleotides specific for said PAP RNA or cDNA suitable for hybridization to PAP RNA or cDNA and/or amplification of PAP sequences and suitable ancillary reagents.

35. A method for increasing a PAP selected from the group consisting of PAP3, PAP7, PAP8, PAP15 and PAP20 in a cell by introducing into said cell a PAP nucleic acid encoding said PAP such that said nucleic acid is expressed and PAP is produced in said cell.

36. A therapeutic method for the treatment or amelioration of diseases resulting from an increase in cell proliferation, said method comprising providing to an individual in need of such treatment an effective amount of an agent or drug which reduces or eliminates PAP expression or function in a pharmaceutically acceptable diluent.

37. The method of claim 36 wherein said disease is cancer.

38. A therapeutic method for the treatment or amelioration of conditions resulting from abnormal cholesterol level, said method comprising providing to an individual in need of such treatment an effective amount of an agent or drug which reduces or eliminates PAP expression or function in a pharmaceutically acceptable diluent.

39. The method of claim 38 wherein said condition is selected from the group consisting of: cancer, neurodegenerative disorders, developmental disorders, stress, and stroke.

40. A method for modulating PBR activity, function or targeting in a cell, said method comprising increasing or decreasing level of PAP in said cell.

FIG. 1A




FIG. 1B




FIG.2A




FIG.2B




FIG.3A





FIG.3B



Tissues and Cell lines

FIG. 4A





FIG. 4B



Tissues and Cell lines

FIG. 5A




FIG. 5B




FIG. 5C




FIG. 5D




FIG. 5E




FIG. 5F



FIG. 5G




FIG. 5H



FIG. 6A




FIG. 6B




FIG. 7A




FIG. 7B




FIG.8A





FIG.8B



SEQ ID NO:1

PAP3 nucleic acid

GAATTCGCGGCCGCGCTCGACACCGCTGCGCCCTCCTGCGAGGCGGGCTGAACGAGGAAATA
ATTGCTAATAAGGCCTCTGTAGCCATGGCTACTTCTGACGTGAAACCAAAATCAATAAGTCGTG
CCAAGAAATGGTCAGAGGAAATAGAAAATCTGTACAGATTTCAACAAGCAGGATATCGGGATG
AAATTGAATATAAACAAGTGAAACAAGTTGCCATGGTCGACCGATGGCCAGAGACAGGGTACG
TGAAGAAACTTTCAGCGGAGGGACAATACTTTCTTCTACTACAACAAAGAGAGGGAGTGCGA
GGACAAGGAGGTCCACAAAGTGAAGGTTTACGTCTACTGACCTTTTCCTTTCTTCGGCTTG
GCAATGCTCCTTTAAGAATTGGTTGTTTACATTCTTCCATCGTGTAATGTCATTTTACAA
ACAATTTCACAATTCTGTCTTTAATTCATGGTGTCTTACACAACATAAACACCCACCTTGA
AACCCAAAA

SEQ ID NO:2

PAP7 sequence:

GAATTCGCGGCCGCGCTCGACCTAAAGTTGAGTTGTTCACTGTAGTGACCCGTGTGAAGGTAGTT
TTATTTTAAATCAACTTTCATTGTGAAACTAGTAAAAGATGGCAAAGCCTTTTCATCCAACCTTA
TGAAGAAAACTGAAGTTCGTGGCACTGCATAAGCAAGTTCTTTTGGGCCCATATAACCCAGA
CACGTCCCCTGAGGTTGGATTCTTTGATGTGTTGGGGAATGATAGGAGGAGAGAATGGGCAGCT
CTGGGAAACATGTCCAAGGAGGATGCCATGGTAGAGTTTGTGAAGCTTCTAAATAAGTGTTGTC
CTCTCCTCTCGGCATATGTTGCGTCCACAGAATAGAGAAGGAAGAAGAAGAGAAAAGAAGAA
AGGCGGAGGAGGAGCGAAGGCAGCGTGAAGAGGAAGAACGAGAGCGGCTGCAAAAGGAAGAA
GAGAAGCGGAAGCGAGAGGAGGAAGACCGGCTGAGACGGGAGGAGGAAGAGAGGGCGGCGGAT
AGAGGAAGAGAGGGCTTCGGCTGGAACAGCAAAAGCAGCAGATAATGGCAGCTTTAAACTCGCA
GACTGCCGTGCAATTCCAGCAGTATGCAGCCAGCAGTATCCAGGGAACCTACGAACAACAGCA
GATTCTCATCCGCCAGCTGCAGGAGCAGCACTATCAGCAGTATAAACACCAGGCAGAGCAAAAC
CCAACCTGCACAACAACAGGCAGCATTACAGAAACAGCAAGAAGTAGTGATGGCTGGGGCAT
CATTGCCTGCATCATCAAAGGTGAACACAGCTGGAGCAAGTGATACACTGTCAGTTAATGGAC
AGGCCAAAACCCACACTGAAAATTCCGAAAAAGTCCTTGAGCCAGAAGCTGCAGAAGAAGCC
TTGGAAAATGGACCAAAAAGACTCTCTTCCAGTGATTGCAGCTCCATCCATGTGGACAAGACCA
CAAATCAAAGACTTTAAAGAGAAGATTCCGCAGGATGCAGATTCTGTGATTACAGTACGTCGA
GGAGAAGTCGTACCGTCCGAGTCCCGACTCATGAGGAAGGATCATACCTATTTTGGGAATTTG
CCACAGACAGTTATGACATTGGGTTTGGGGTTTATTTGAATGGACAGACTCTCCAAATGCTGC
TGTCAGTGTGCATGTCAGTGAGTCCAGTGACGAGGAGGAGGAGGAGGAAGAAAATGTCACTTG
TGAAGAAAAAGCAAAAAAGAACGCCAACAAGCCTCTGCTGGATGAGATTGTACCTGTGTACCG
GCGGACTGTACGAGGAAGTATATGCAGGCAGCCACCAGTATCCAGGGAGGGGAGTCTATC
TCCTCAAGTTTGATAATTCTTACTCTCTGTGGAGGTCCAAGTCCGTCTACTACAGAGTCTA
TTATACTAGATAGAGCTGCTGTTCCAAGGTCCGGAGTCCAGGGTTGAGCACAACATGACGT
TTAATTTCTCTT

SEQ ID NO:3

PAP8 nucleic acid

GAATTCGCGGCCGCGCTCGACGCTGGACACAAGCGTGGAGCGAAGAGCCCTTGGAGAGATTTCAG
AATGTAGGTGAAGGCTCTTCAACATCACAGGGCACCTGGCAGTCTTCAGAGTCCTCACAGTCA
AACCTGGGGGAGCAGACGCAGAGCGGACCCAGGGAGGAAGGTGTCAGCGTCGGGAGAGGGCA
TAACCGAATGGAACGAGATAGGAGGCGCAGAATCCGCATTTGCTGTGATGAGCTGAATCTTTTA
GTTCCATTCTGCAATGCGGAGACAGATAAAGCAACAACCCTTCAGTGGACCACAGCATTCCTG
AAGTACATTTCAGGAAAGACATGGGGACTCTCTTAAAAAGGAATTTGAGAGCGTGTTTTGCGGTA
AAACTGGCAGAAGGCTAAAGCTGACTAGACCCGAATCCCTGGTGACCTGCCCTGCACAGGGCA
GCCTGCAGAGCAGCCCTGCCATGGAGATCAAGTGACTGGACTGACCCAGGACCTGGGAGAGA

ACCGCCGTTCTCGGGCATCATGCACATGCCTGCCATCCCCGGAATTCAGCTCTGAATCCT
CTC

SEQ ID NO:4

PAP15 nucleic acid

GAATTCGCGGCCGCGTCGACCTCGGGGGTAACCCTTGGGATCATTATGATGTCACCTCTTCAAT
CTCTTGAGACCAGTGATTTATGAACTTGACATTTGGTGCCTGGTGTGGGCTTTGGAAAGCAGAG
GCCACCTTTGTCTGTAGAGGATACTGAGCGGCTGGATGGCAGGAATCCAAAGAGAAGCCAGCC
TCTCGTAGTCGCCCTGGGACAGTGGAAGAGAGGGAGGACCGGCAAAGGGGAATCTGTCTTTCT
CCTAGGCCCCGAGCATGTCCCCTGTGGGACATGCTCTGTGACAGCTGAGCCTGCCCAGCCTG
CCTTTCTGAAGTTGGGTGTCTCCTGCCCACAACCAAGCCAGCAATCGGTCTGTTTTCCGAC
AACCTCAGAGCCAGACCTCACAAAGCCTATTTTGGTGGTTCCCAAATTTCTCTCAGATCTC
CATGTCTATCCCTCCACTCCCTCCAAAGAGAAAGAAAGAATTGAGAAAGAA

SEQ ID NO:5

PAP20 nucleic acid

GAATTCGCGGCCGCGTCGACCTCAGAAGAGGAAAAGAGGTGCAGAAGTGCTGGCGGCACAAA
TTGTACAGAAAACCAGACTAGAGAGAAAAAACAAGAAGCGTCTGTATCTAAAGATGCTCCAG
TGCCTACAAATACTAAAAGGGCAAAGAAACAAGAGAAGTCTCCAGGTAGAATTGCCTCACAGT
CTAAGCCACCCATGAAAAAGTCTCCACAAAAACGGAAGGTAAATGTAGCAAGAGGCCGTCGG
AATACCAGAAAGCAGCTCCAACCTGCCGAAAAAGAAATTGCTTTACATCTTCAATCAGAAATT
TCATCAGATGGCCAAAAAGATGGACTTAACCTAAGTACATCTCAACAAGAAAGTATTTCAATG
ATTCTAAAGGTCTCCTGAAAACCTCAGTTATCAGCTGTGACTCCCAGGCCCTAAATATGTTAG
CTGATCTGGCATTAAAGTTCTGCTGCTGCTTCTATACCATCTTGTAAGCCCAGGAACCTTCCCTG
CGTCTCTGATTTGCCACGAAACAATGTCTTACTCACTAAAGAAAATCCATTGCTTGGTGCCT
CTGACCATGAATATCATAAGGG

SEQ ID NO:6

PAP3 amino acid sequence

Met Ala Thr Ser Asp Val Lys Pro Lys Ser Ile Ser Arg Ala Lys
Lys Trp Ser Glu Glu Ile Glu Asn Leu Tyr Arg Phe Gln Gln Ala
Gly Tyr Arg Asp Glu Ile Glu Tyr Lys Gln Val Lys Gln Val Ala
Met Val Asp Arg Trp Pro Glu Thr Gly Tyr Val Lys Lys Leu Gln
Arg Arg Asp Asn Thr Phe Phe Tyr Tyr Asn Lys Glu Arg Glu Cys
Glu Asp Lys Glu Val His Lys Val Lys Val Tyr Val Tyr

SEQ ID NO:7

PAP7 amino acid sequence

Arg Pro Arg Arg Pro Lys Val Glu Leu Phe Thr Val Val Thr Arg
Val Lys Val Val Leu Phe Leu Asn Gln Leu Ser Leu Cys Lys Leu
Val Lys Asp Gly Lys Ala Phe His Pro Thr Tyr Glu Glu Lys Leu
Lys Phe Val Ala Leu His Lys Gln Val Leu Leu Gly Pro Tyr Asn
Pro Asp Thr Ser Pro Glu Val Gly Phe Phe Asp Val Leu Gly Asn
Asp Arg Arg Arg Glu Trp Ala Ala Leu Gly Asn Met Ser Lys Glu
Asp Ala Met Val Glu Phe Val Lys Leu Leu Asn Lys Cys Cys Pro

Leu Leu Ser Ala Tyr Val Ala Ser His Arg Ile Glu Lys Glu Glu
 Glu Glu Lys Arg Arg Lys Ala Glu Glu Arg Arg Gln Arg Glu
 Glu Glu Glu Arg Glu Arg Leu Gln Lys Glu Glu Glu Lys Arg Lys
 Arg Glu Glu Glu Asp Arg Leu Arg Arg Glu Glu Glu Glu Arg Arg
 Arg Ile Glu Glu Glu Arg Leu Arg Leu Glu Gln Gln Lys Gln Gln
 Ile Met Ala Ala Leu Asn Ser Gln Thr Ala Val Gln Phe Gln Gln
 Tyr Ala Ala Gln Gln Tyr Pro Gly Asn Tyr Glu Gln Gln Gln Ile
 Leu Ile Arg Gln Leu Gln Glu Gln His Tyr Gln Gln Tyr Lys His
 Gln Ala Glu Gln Thr Gln Pro Ala Gln Gln Gln Ala Ala Leu Gln
 Lys Gln Gln Glu Val Val Met Ala Gly Ala Ser Leu Pro Ala Ser
 Ser Lys Val Asn Thr Ala Gly Ala Ser Asp Thr Leu Ser Val Asn
 Gly Gln Ala Lys Thr His Thr Glu Asn Ser Glu Lys Val Leu Glu
 Pro Glu Ala Ala Glu Glu Ala Leu Glu Asn Gly Pro Lys Asp Ser
 Leu Pro Val Ile Ala Ala Pro Ser Met Trp Thr Arg Pro Gln Ile
 Lys Asp Phe Lys Glu Lys Ile Arg Gln Asp Ala Asp Ser Val Ile
 Thr Val Arg Arg Gly Glu Val Val Thr Val Arg Val Pro Thr His
 Glu Glu Gly Ser Tyr Leu Phe Trp Glu Phe Ala Thr Asp Ser Tyr
 Asp Ile Gly Phe Gly Val Tyr Phe Glu Trp Thr Asp Ser Pro Asn
 Ala Ala Val Ser Val His Val Ser Glu Ser Ser Asp Glu Glu Glu
 Glu Glu Glu Glu Asn Val Thr Cys Glu Glu Lys Ala Lys Lys Asn
 Ala Asn Lys Pro Leu Leu Asp Glu Ile Val Pro Val Tyr Arg Arg
 Asp Cys His Glu Glu Val Tyr Ala Gly Ser His Gln Tyr Pro Gly
 Arg Gly Val Tyr Leu Leu Lys Phe Asp Asn Ser Tyr Ser Leu Trp
 Arg Ser Lys Ser Val Tyr Tyr Arg Val Tyr Tyr Thr Arg

SEQ ID NO:8

PAP8 amino acid sequence

Glu Phe Ala Ala Ala Ser Thr Leu Asp Thr Ser Val Glu Arg Arg
 Ala Leu Gly Glu Ile Gln Asn Val Gly Glu Gly Ser Ser Thr Ser
 Gln Gly Thr Trp Gln Ser Ser Glu Ser Ser Gln Ser Asn Leu Gly
 Glu Gln Thr Gln Ser Gly Pro Gln Gly Gly Arg Cys Gln Arg Arg
 Glu Arg His Asn Arg Met Glu Arg Asp Arg Arg Arg Arg Ile Arg
 Ile Cys Cys Asp Glu Leu Asn Leu Leu Val Pro Phe Cys Asn Ala
 Glu Thr Asp Lys Ala Thr Thr Leu Gln Trp Thr Thr Ala Phe Leu
 Lys Tyr Ile Gln Glu Arg His Gly Asp Ser Leu Lys Lys Glu Phe
 Glu Ser Val Phe Cys Gly Lys Thr Gly Arg Arg Leu Lys Leu Thr
 Arg Pro Glu Ser Leu Val Thr Cys Pro Ala Gln Gly Ser Leu Gln
 Ser Ser Pro Ala Met Glu Ile Lys

SEQ ID NO:9

PAP15 amino acid sequence

Ala Ala Gly Trp Gln Glu Ser Lys Glu Lys Pro Ala Ser Arg Ser
 Arg Pro Gly Thr Val Glu Glu Arg Glu Asp Arg Gln Arg Gly Ile
 Cys Leu Ser Pro Arg Pro Glu His Val Pro Cys Gly Thr Cys Ser
 Val Thr Ala Glu Pro Ala Gln Pro Ala Phe Leu Lys Leu Gly Val
 Ser Cys Pro Gln Pro Ser Gln Gln Ser Val Cys Phe Pro Thr Thr
 Ser Glu Pro Asp Leu Thr Ser Leu Phe Trp Trp Phe Pro Lys Phe
 Leu Ser Asp Leu His Val Tyr Pro Ser Thr Pro Ser Lys Arg Glu
 Arg Lys Glu Leu Arg Lys Lys

SEQ ID NO:10

PAP20 amino acid sequence

Asn Ser Arg Pro Arg Arg Pro Gln Lys Arg Lys Arg Gly Ala Glu
Val Leu Ala Ala Gln Ile Val Gln Lys Thr Arg Leu Glu Arg Lys
Lys Gln Glu Ala Ser Val Ser Lys Asp Ala Pro Val Pro Thr Asn
Thr Lys Arg Ala Lys Lys Gln Glu Lys Ser Pro Gly Arg Ile Ala
Ser Gln Ser Lys Pro Pro Met Lys Lys Ser Pro Gln Lys Arg Lys
Val Asn Val Ala Arg Gly Arg Arg Asn Thr Arg Lys Gln Leu Gln
Pro Ala Glu Lys Glu Ile Ala Leu His Leu Gln Ser Glu Ile Ser
Ser Asp Gly Gln Lys Asp Gly Leu Asn Leu Ser Thr Ser Gln Gln
Glu Ser Ile Ser Met Ile Pro Lys Gly Pro Pro Glu Asn Ser Val
Ile Ser Cys Asp Ser Gln Ala Leu Asn Met Leu Ala Asp Leu Ala
Leu Ser Ser Ala Ala Ala Ser Ile Pro Ser Cys Lys Pro Arg Asn
Leu Pro Cys Val Ser Asp Leu Pro Arg Asn Asn Val Leu Leu Thr
Lys Glu Asn Pro Leu Leu Gly Ala Ser Asp His Glu Tyr His Lys
Gly

SEQ ID NO:11

Ser Ser Asp Glu Glu Glu Glu Glu Glu Glu Asn Val Thr Cys Glu
Glu Lys Ala Lys Lys Asn Ala Asn Lys Pro



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/12, C07K 14/47, C12N 1/21, G01N 33/53, C07K 16/18, C12Q 1/68, A61P 35/00, 3/06	A3	(11) International Publication Number: WO 00/09549 (43) International Publication Date: 24 February 2000 (24.02.00)
(21) International Application Number: PCT/US99/18507 (22) International Filing Date: 11 August 1999 (11.08.99) (30) Priority Data: 60/096,048 11 August 1998 (11.08.98) US (71) Applicant (for all designated States except US): GEORGE-TOWN UNIVERSITY MEDICAL CENTER [US/US]; Office of the Dean of Research and Graduate Education, Building D, suite 177, Washington, DC 20007 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): PAPADOPOULOS, Vasilios [US/US]; 15417 Peach Leaf Drive, North Potomac, MD 20878 (US). LI, Hua [CN/US]; Apartment 102, 1319 Fort Myer Drive, Arlington, VA (US). (74) Agent: PRATT, Sana, A.; Pratt & Associates, Inc., 10821 Hillbrooke Lane, Potomac, MD 20854 (US).		(81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> (88) Date of publication of the international search report: 16 November 2000 (16.11.00)
(54) Title: PERIPHERAL-TYPE BENZODIAZEPINE RECEPTOR ASSOCIATED PROTEINS, CLONING, EXPRESSION AND METHODS OF USE (57) Abstract The present invention relates to nucleic acids encoding PBR-associated proteins (PAPs) and methods for use in producing PAPs and methods of using PAPs. In this study we have identified proteins (PAPs) that interact with PBR protein by a yeast two-hybrid system. We used PBR as bait to screen a mouse testis cDNA library. Five clones were isolated based on their ability to interact with PBR. These proteins may be involved in the regulation of the function of PBR, serving as an endogenous ligand or allosteric modulator of the receptor.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/US 99/18507

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C07K14/47 C12N1/21 G01N33/53 C07K16/18
C12Q1/68 A61P35/00 A61P3/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K G01N C12Q A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL 'Online! AC AF022770, 1 October 1997 (1997-10-01) LI H & V PAPADOPOULOS: "M. musculus peripheral benzodiazepine receptor associated protein" XP002133903 cited in the application 94.9% identity in 178 aa overlap with SEQ ID No:7 the whole document</p> <p style="text-align: center;">--- -/-</p>	<p>1-5, 9-16,19</p>

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

15 June 2000

Date of mailing of the international search report

28.06.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040. Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Nichogiannopoulou, A

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 99/18507

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE EMBL 'Online! AC X64455, 21 February 1992 (1992-02-21) DON J: "M. musculus mRNA meg1" XP002133904 cited in the application 99% identity in 490 nt overlap with SEQ ID No:1 (17-505:28-517) the whole document</p> <p>---</p>	3,9-16
X	<p>DATABASE EMBL 'Online! AC AA174581, 31 December 1996 (1996-12-31) MARRA M ET AL: "mo98e12.r1 Stratagene mouse testis (#937308) Mus musculus cDNA" XP002133905 EST with 97.9% identity in 421 nt overlap (17-435:79-498) with SEQ ID No:1 the whole document</p> <p>---</p>	3,9-16
E	<p>WO 00 01821 A (INCYTE PHARMA INC) 13 January 2000 (2000-01-13) Incyte clone with 88.7% identity to SEQ ID NO:2 table 2</p> <p>---</p>	3-5, 9-16,19
P,A	<p>GALIÈGUE S ET AL: "Cloning and characterisation of PRAX-1. A new protein that specifically interacts with the peripheral benzodiazepine receptor" J. BIOL. CHEM, vol. 274, no. 5, 29 January 1999 (1999-01-29), page 2938-2952 XP002122735 cited in the application the whole document</p> <p>---</p>	
A	<p>KRUEGER K E ET AL: "Peripheral-type benzodiazepine receptors mediate translocation of cholesterol from outer to inner mitochondrial membranes in adrenocortical cells" JOURNAL OF BIOLOGICAL CHEMISTRY., vol. 265, no. 25, 5 September 1990 (1990-09-05), pages 15015-15022, XP002133901 ISSN: 0021-9258 cited in the application the whole document</p> <p>---</p>	38,39

	-/--	

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/US 99/18507

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	MIETTINEN H ET AL: "Expression of peripheral-type benzodiazepine receptor and diazepam binding inhibitor in human astrocytomas: relationship to cell proliferation" CANCER RESEARCH, vol. 55, 15 June 1995 (1995-06-15), pages 2691-2695, XP002133902 cited in the application the whole document -----	36,37

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 99/18507

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 35-40 -in as far as they concern in vivo methods- are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: 29-32
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

The claimed agents have not been described in the application and have as such not been the subject of search.
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☒ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:

1-4, 9-18, 23-25, 27-40 partially and 5, 19, 26 completely
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 29-32

The claimed agents have not been described in the application and have as such not been the subject of search.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-3, 9-18, 24, 25, 27-32, 35-40 all partially

An isolated DNA fragment encoding PBR-associated protein 3 (PAP3) and portions of said DNA, including the DNA of SEQ ID No:1 and fragments of said sequence of at least 30 nucleotides. Recombinant DNA constructs, host cells and production methods relating to PAP3. Recombinant PAP3 and detection methods comprising the two hybrid assay. Antibodies to the PAP3 polypeptide of SEQ ID No:6. Methods for detecting agents which reduce, eliminate or promote PAP3 activity, agents and therapeutic compounds that inhibit or promote PAP activity and therapeutic methods for the treatment of disease states.

2. Claims: 5, 19, 26 completely and 1-4, 9-18, 23-25, 27-40 all partially

As invention 1 but relating to PAP7 (SEQ ID Nos: 2 and 7)

3. Claims: 6, 20 completely and 1-3, 9-18, 23-25, 27-40 all partially

As invention 1 but relating to PAP8 (SEQ ID Nos: 3 and 8)

4. Claims: 7, 21 completely and 1-3, 9-18, 23-25, 27-40 all partially

As invention 1 but relating to PAP15 (SEQ ID Nos: 4 and 9)

5. Claims: 8, 22 completely and 1-4, 9-18, 23, 24, 27-40 all partially

As invention 1 but relating to PAP20 (SEQ ID Nos: 5 and 10)

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/US 99/18507

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0001821 A	13-01-2000	AU 4857999 A	24-01-2000